

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89309257.7

(51) Int. Cl.⁵: **A 61 K 37/02**
C 07 K 7/64

(22) Date of filing: 12.09.89

(30) Priority: 12.09.88 US 242767 16.09.88 US 246042
30.08.89 US 398889

(43) Date of publication of application:
21.03.90 Bulletin 90/12

(84) Designated Contracting States:
AT BE CH DE ES FR GB IT LI LU NL SE

(71) Applicant: MERCK & CO. INC.
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900 (US)

(72) Inventor: Schmatz, Dennis M.
306 Elizabeth Avenue
Cranford New Jersey 07016 (US)


(74) Representative: Hesketh, Alan, Dr. et al
European Patent Department Merck & Co., Inc. Terlings
Park Eastwick Road
Harlow Essex, CM20 2QR (GB)

The applicant has filed a statement in accordance with Rule 28 (4) EPC (Issue of a sample only to an expert). Accession number(s) of the deposit(s): ATCC 20868 and ATCC 20956.
The microorganism(s) has (have) been deposited with American Type Culture Collection under number(s) ATCC 20868 and ATCC 20956.

(54) Method for the control of pneumocystis carinii.

(57) A method for the treatment of *Pneumocystis carinii*, the causative agent of pneumonia of particular severity to immune compromised patients such as those with acquired immune deficiency syndrome (AIDS) by administering a lipophilic cyclohexapeptide compound is described. Compositions suitable for the treatment of *P. carinii* are also disclosed.

FIG-1



Description

METHOD FOR THE CONTROL OF PNEUMOCYSTIS CARINII

BACKGROUND OF THE INVENTION

Pneumocystis carinii is an opportunistic organism which is widely prevalent but generally dormant until bodily defenses of the host are compromised whereupon it propagates in the host causing disease. Generally it is present in the lungs although extrapulmonary infections have been reported. Lung infections with this organism in immunocompromised individuals generally lead to pneumonia and are usually fatal if untreated. It has been suggested that patient to patient transmission also may occur in immunocompromised individuals. The immunocompromised state of individuals is usually associated with genetic defects, lymphoproliferative diseases, or with conditions resulting from cancer therapy, or treatment with immunosuppressive drugs, or as a consequence of AIDS. Most AIDS patients eventually contract *Pneumocystis carinii* pneumonia (PCP) and have accounted for the majority of the recent cases of this disease. Left untreated, PCP is almost always fatal.


The current method of treatment for *P. carinii* pneumonia is trimethoprim/sulfamethoxazole or pentamidine. Treatment with trimethoprim/sulfamethoxazole (TMP/SMZ) is associated with a high level of toxic side effects including rash, elevated liver function, nausea and vomiting, anemia, creatine elevation, and in extreme cases Stevens-Johnson syndrome. Side effects from TMP/SMZ are much more prevalent in patients with AIDS. Treatment with pentamidine is also associated with a high level of toxic side effects including renal failure, hepatotoxicity, hypoglycemia, hematologic abnormalities and pain or abscess at the injection site. The mortality due to treatment can reach 20 to 30 percent. An improved method for the treatment of *P. carinii* pneumonia in immune-compromised patients is greatly needed.

DESCRIPTION OF THE INVENTION

The present invention is directed to a method for treatment or prevention of *Pneumocystis carinii* infections in mammals which comprises administering to subjects, a person or animal, infected with or susceptible to being infected with *Pneumocystis carinii*, an anti-infection or infection controlling amount of a cyclohexapeptide compound.

The compounds within the scope of the present invention include a novel fungal metabolite as hereinafter detailed, known echinocandin type of antibiotics which are lipophilic cyclohexapeptides having a fatty acid side chain, simple derivatives of the echinocandin type antibiotics such as side chain dehydro- and tetrahydro-reduction products, ring ether and ester derivatives and semi-synthetic products in which the cyclohexapeptide ring is retained but in which the fatty acid side chain is replaced. Most of the compounds described in the literature are taught to be antifungal agents.

The compounds suitable for the practice of the present invention which include not only natural antibiotic compounds but also simple derivatives of natural antibiotic compounds and semi-synthetic compounds as are hereinafter fully detailed may be represented by the formula (I).




In this and succeeding formulas

R' is H or OH;

R'' is H or OH;

R''' is H, OH, -O-alkyl(C₁-C₆), -O-benzyl, or



wherein R¹ is H, C₁-C₁₂ alkyl, C₃-C₇ cycloalkyl, hydroxyethyl, lower alkoxyethyl, furylmethyl, tetrahydrofurylmethyl, phenyl, phenylalkyl wherein the phenyl may be substituted with halo, hydroxy, lower alkyl and lower alkoxy; R² is hydrogen or lower alkyl; or R¹ and R² together form -(CH₂)₂Q(CH₂)₂- wherein Q is a connecting group such as -CH₂-, -NH- or -N-(lower alkyl)-; and m is from 0 to 4;

R''' is H or OH;



R''' is H, CH₃ or CH₂CONH₂; and

R''' is H or CH₃


R is (A) -C(=O)-alkyl, preferably from 6 to 24 carbon atoms, or

(B) -C(=O)-alkenyl, preferably from 6 to 24 carbon atoms,

(C) when R', R'', R''' and R''' are OH, and R''' and R''' are CH₃ then



or



wherein

A is divalent oxygen, sulfur, sulfinyl or sulfonyl;

A' is divalent oxygen, sulfur, sulfinyl, sulfonyl or -NH-;

X is hydrogen, chloro, bromo, iodo, nitro, C₁-C₃ alkyl, hydroxy, C₁-C₃ alkoxy, mercapto, C₁, C₃ alkylthio, carbamyl or C₁-C₃ alkylcarbamyl;

X' is chloro, bromo or iodo;

R₁ is hydrogen, C₁-C₁₈ alkyl or C₂-C₁₈ alkenyl;

W is C₁-C₁₀ alkylene or C₂-C₁₀ alkenylene;


m, n and p are 0 or 1, but if m is 0, n must be 0;

provided that the sum of the carbon atoms in the R₁ and W groups must be greater than 4 but cannot exceed 21; that when X is mercapto A and A' cannot be sulfinyl or sulfonyl; and that when A and A' are sulfinyl or sulfonyl, they must be in equal oxidation states;


(D) when R' is OH, R'', R''' and R''' are H, then



wherein Y is a divalent aminoacyl radical of the formula




wherein Z is C₁-C₁₀ alkylene or C₅-C₆ cycloalkylene



wherein R₃ is hydroxymethyl, hydroxyethyl, mercaptomethyl, mercaptoethyl, methylthioethyl, 2-thienyl, 3-indolomethyl, phenyl, benzyl, halophenyl, halobenzyl, C₁-C₃ alkylphenyl, C₁-C₃ alkylbenzyl, C₁-C₃

alkylthiophenyl, C₁-C₃ alkylthiobenzyl, carbamylphenyl, carbamylbenzyl, C₁-C₃ alkylcarbamylphenyl or C₁-C₃ alkylcarbamylbenzyl;
(c)




10 wherein X² is hydrogen or halo.


The term "alkyl" is meant a univalent saturated straight chain or branched chain hydrocarbon radical. The term "alkenyl" is meant a univalent, unsaturated, straight chain or branched chain hydrocarbon radical containing not more than three double bonds. By "lower" is meant up to 6 carbon atoms. The term "halo" is meant chloro, bromo, fluoro and iodo.

15 The compounds useful in the method of treatment for or prevention of *Pneumocystis carinii* infections in mammals according to the present invention are generally white crystalline or powdery substances. They are generally of good solubility in alcohols such as methanol and ethanol, or poor solubility in water and in hydrocarbons such as hexane and of intermediate solubility in many common solvents. The solubility properties as well as other properties of many of the naturally occurring cyclohexapeptides are summarized in the CRC Handbook of Antibiotic Compounds, Vol IV, Part I, pp 355-367, CRC Press, Inc. Boca Raton, Florida, 1980.

An especially preferred compound is a novel fungal metabolite in which R', R'', R''' and R'''' are OH, R'''' is -CH₂CONH₂, R''' is CH₃ and R is



30 represented by formula IA



55 The compound may be named 1-[4,5-dihydroxy-N²-(10,12-dimethyl-1-oxotetradecyl)-L-ornithine-5-(threo-3-hydroxy-1-glutamine)echinocandin B but hereinafter, for convenience, will be referred to as Compound IA. Especially preferred is the isomer named 1-[(4R,5R)-4,5-dihydroxy-N²-(10,12-dimethyl-1-oxo-tetradecyl)-L-ornithine-5-(threo-3-hydroxy-1-glutamine)echinocandin B. The preparation and properties of Compound IA are hereinafter described and are also the subject of copending applications Serial No. 105,795, filed October 7, 1987, corresponding to EP 311193, and Serial No. 105,797, filed October 7, 1987, corresponding to EP 311194, and are more fully described therein.

60 Compound IA, the preferred compound, is a white solid which may be characterized by the following physical properties:

a decomposition temperature of 206-214°C;

an empirical formula of C₅₁H₈₂N₈O₁₇ determined by high resolution FAB (Fast Atom Bombardment mass spectrometric measurement, calculated for C₅₁H₈₂N₈O₁₇ + Li = 1085.5958, found = 1085.6146);

an amino acid composition as determined by gas chromatogram/mass spectra of the trimethylsilyl derivative of the total acid hydrolysates of one equivalent each of threonine, hydroxyproline, methylhydroxyproline, and hydroxyglutamic acid.

¹H NMR Spectra in CD₃OD at 400 MHz as seen in Figure 1; and

¹³C NMR chemical shifts obtained in CD₃OD at 100 MHz as follows: 11.20, 11.64, 19.75, 20.25, 20.78, 27.00, 28.07, 30.33, 30.37, 30.61, 30.76, 31.19, 31.29, 32.94, 34.83, 36.69, 38.10, 38.54, 39.07, 39.53, 45.93, 51.39, 53.01, 55.59, 56.31, 57.11, 58.35, 62.43, 68.18, 70.08, 70.55, 70.61, 71.26, 73.94, 75.72, 75.84, 76.86, 116.06(x2), 129.43(x2), 132.86, 158.22, 168.80, 172.16, 172.35, 172.40, 173.12, 174.24, 175.47, 176.88 ppm.

The compound is soluble in a variety of organic solvent such as methanol, ethanol, dimethylformamide, dimethyl sulfoxide, and the like.

The production of Compound IA, a novel compound, is described hereinafter in detail. The other compounds useful in the present invention may be prepared in a similar manner by substituting an appropriate microorganism or by other procedures as subsequently described.

Compound IA may be produced by cultivating a strain of microorganism designated MF 5171 in the culture collection of Merck & Co., Rahway, N.J. and identified as *Zalerion arboricola* and recovering said agent from the culture medium. A sample of the culture capable of producing the Compound IA has been deposited under the Budapest Treaty in the Culture Collection of the American Type Culture Collection at 12301 Parklawn Drive, Rockville, MD 20852. The sample has been assigned the accession number ATCC 20868.

The colonial and morphological description of ATCC 20868 are set forth below:

A. Morphological description

Globose, approximately 6.0 microns in diameter, thick-walled, dark coloured structures similar to chlamydospores develop along mycelium and often appear to be intercalary. These structures appear to divide, forming multi-celled groups of 4-8 cells which do not break up readily. On some media, strands of mycelia cluster together, forming rope-like structures and the multi-celled groups form large clusters as if held together by mucilaginous material.

B. Colonial description

1. Czapek-Dox agar

Colonies are slow growing, growth not extensive, flat with irregular edges. Mycelium is black, shiny; surface becomes dull, granular-appearing, black to greenish-brown as colony ages.

2. Corn agar

Colonies are slow-growing. Growth is not extensive. Mycelium is black, shiny surface becoming powdery, dull black as colony ages.

3. Potato-dextrose agar, Sabouraud maltose agar and yeast extract-malt extract-dextrose agar.

Colonies are slow growing. Growth is not extensive, is slightly raised in the center, radiately furrowed irregular edges except on Sabouraud-maltose agar where the edges are hyaline and regular. Mycelium is black, shiny; becomes dull powder, black as colony ages.

Although the invention is discussed herein below principally with respect to the specific strain, it is well-known in the art that the properties of microorganisms may be varied naturally and artificially. Thus, all strains of the genus ATCC 20868 including varieties and mutants, whether obtained by natural selection, produced by the action of mutating agents such as ionizing radiation or ultraviolet irradiation, or by the action of chemical mutagens such as nitrosoguanidine, are contemplated to be within the scope of this invention.

Compound IA may be produced in a form adaptable for drug use by cultivating the strain ATCC 20868 of *Zalerion arboricola* in a nutrient medium until a substantial amount of antibiotic activity is detected in the culture medium, generally in the mycelium portion thereof, and thereafter recovering the active component from the fermentation mycelium in a suitable solvent, concentrating the solution of active component, and then subjected the concentrated material to chromatographic separation.

The fermentation is carried out in a medium containing sources of carbon and nitrogen assimilable by the microorganisms and generally low levels of inorganic salts. In addition, the medium may be supplemented with trace metals, although if complex sources of carbon and nitrogen are employed, they are usually present in the complex sources.

The source of carbon include glycerol, sugars, starches and other carbohydrates or carbohydrate derivatives such as dextran, cellose, as well as complex nutrients such as oat flour, corn meal, millet, corn and the like. The exact quantity of the carbon source which is utilized in the medium will depend, in part, upon the other ingredients in the medium, but it is usually found that an amount of carbohydrate between 0.5 and 5 percent by weight of the medium is satisfactory. These carbon sources can be used individually or several such carbon sources may be combined in the same medium.

The sources of nitrogen include amino acids such as glycine, arginine, threonine, methionine and the like as well as complex sources such as yeast hydrolysates, yeast autolysates, yeast cells, tomato paste, soybean meal, casein hydrolysates, yeast extracts, corn steep liquors, distillers solubles, cottonseed meal, meat extract, and the like. The various sources of nitrogen can be used alone or in combination in amounts ranging from 0.2 to 90 percent by weight of the medium.

Among the nutrient inorganic salts, which can be incorporated in the culture media are the customary salts

EP 0 359 529 A1

capable of yielding sodium, potassium, magnesium, ammonium, calcium, phosphate, sulfate, chloride, carbonate, and like ions. Also included are trace metals such as cobalt, manganese, iron, molybdenum, zinc, cadmium, and the like.

The media suitable for carrying out the fermentation may be solid or liquid.

- 5 Solid medium may have a millet, corn, oats, soy bean or wheat base. One medium having a millet base is Medium 2 in Example A. Other representative solid media include the following:

10	Media	Weight or Volume Per 250 ml Flask
15	<u>Medium A</u>	
	Corn (cracked)	10.0 g
	Yeast extract	0.5 g
20	Sodium tartrate	0.1 g
	Monosodium glutamate	0.1 g
	Corn oil	0.1 ml
25	Ferrous sulfate•7H ₂ O	0.01 g
	Water	15-20 ml
	<u>Medium B</u>	
30	Millet	15 g
	Yeast extract	0.5 g
	Sodium tartrate	0.1 g
35	Ferric sulfate•7H ₂ O	0.01 g
	Sucrose	0.5 g
	Alfalfa	0.5 g
40	Corn oil	0.1 ml
	Water	15 ml
45	<u>Medium C</u>	
	Millet	15 g
	Yeast extract	0.5 g
50	Sodium tartrate	0.1 g
	Ferric sulfate•7H ₂ O	0.01 g
	Silica gel	0.5 g
55	Alfalfa	0.5 g
	Monosodium glutamate	0.1 g
	Corn oil	0.1 ml
60	Water	15 ml

65

Medium D

Millet	15 g	
Yeast extract	0.5 g	5
Sodium tartrate	0.1 g	
Ferric sulfate•7H ₂ O	0.01 g	10

In addition, media may be prepared by substituting wheat, barley, oats or soy bean for the millet or corn above.

Liquid media also may be employed. Fermentation on a larger scale is generally more conveniently carried out employing a liquid medium. It has been found that although conventional liquid media may be employed to obtain Compound IA, such media have not been found to be suitable for obtaining good yields of the desired antibiotic. However, by incorporating from about 6 to 9 percent by weight of glycerol, it has been found that good yields of the desired antibiotic compound may be obtained. Suitable liquid media include:

Medium E

Glycerol	85 g	20
Pectin	10 g	
Peanut Meal	4 g	25
Peptonized Milk	4 g	
Tomato Paste	4 g	
Corn Steep	4 g	30
Lard Water	4 g	
Glycine	2 g	
KH ₂ PO ₄	2 g	35
Distilled Water to	1000 ml	

pH 7.0

Medium F

Dextrose	10 g	
Glycerol	10 ml	50
Soy Flour	4 g	
Peptonized milk	4 g	55
Tomato paste	4 g	
Lard water	4 g	
Potassium dihydrogen phosphate	2 g	60
Cobalt chloride hexahydrate	0.01 g	
Distilled Water to	1000 ml	

pH 7.0

For producing Compound IA, a fermentation medium containing ATCC 20868 is prepared by inoculating spores or mycelia of the antibiotic-producing organism into a suitable medium and then cultivating under aerobic conditions.

5 The procedure generally is first to inoculate a preserved source of culture from an agar slant containing nutrient medium into a nutrient seed-producing medium and to obtain, preferably through a two step procedure, growth of the organisms which serve as seeds in the production of the antipneumocystis agent.

In this process, a slant section of a preserved culture of ATCC 20868 is inoculated into an appropriate liquid nutrient seed medium of pH in the range 5 to 8.1, optimally 6 to 7.5, and the flasks incubated with or without agitation at temperatures in the range of from about 15°C to about 30°C, preferably 20° to 28°C. Agitation when employed, may be up to 400 rpm, preferably, about 200 to 220 rpm. The incubation is carried out over a period of from 2 to 30 days, preferably 2 to 14 days. When the growth is abundant, usually between 2 and 5 days, the culture growth may be used to inoculate the production medium for the production of the antipneumocystis agent. Preferably however, a second stage fermentation is carried out, inoculating with a portion of the culture growth and then employing similar conditions but generally with shortened incubation period of about 1 to 3 days. The growth then is employed to inoculate the production medium.

The fermentation production medium inoculated with the culture growth is incubated for 3 to 30 days, usually 7 to 14 days with or without agitation. The fermentation may be conducted aerobically at temperatures ranging from about 20°C to about 40°C. For optimum results, it is most convenient to conduct these fermentations at a temperature in the range of from about 24°C to about 30°C. Temperatures of about 24°C-28°C are most preferred. The pH of the nutrient medium suitable for producing the instant compounds can vary from about 5.0 to 8.5 with a preferred range of from about 6.0 to 7.5. After the appropriate period for the production of the desired compound or compounds, the latter is recovered from the fermentation medium as hereinafter more fully described.

25 The active material may be recovered from the fermentation medium by the steps comprising

- (1) adding alcohol to said medium, stirring and filtering to recover the active component in the resulting aqueous alcoholic solution;
- (2) concentrating the aqueous alcoholic solution to a small volume of primarily aqueous solution;
- (3) intimately contacting the resulting concentrated alcoholic aqueous solution with a water-immiscible organic solvent to extract or partition the active component thereinto and concentrating, or subjecting the concentrated solution to HP-20 adsorption and elution; then
- (4) subjecting the material recovered in Step (3) to at least one chromatographic separation, wherein in each chromatographic separation, the active component from the eluates exhibiting activity against Candida albicans are combined and concentrated to recover Compound IA.

35 The exact steps may vary somewhat depending on whether the fermentation had been carried out in liquid or solid medium, what solvent is employed and what adsorbent or combination of adsorbents is employed.

When the fermentation is carried out on solid medium, the first step may be carried out by adding an alcoholic solvent to the fermentation medium, thoroughly mixing, then filtering, recovering and concentrating the aqueous alcohol filtrate. Preferably, the concentrated filtrate is first back-extracted or washed with a lower aliphatic hydrocarbon solvent such as hexane or other alkane to remove alkane soluble impurities. The alkane washed filtrate may be extracted or partitioned with a water-immiscible oxygenated organic solvent and the resulting solution concentrated, then placed onto a column for at least one, generally several chromatographic separation steps. Suitable columns are silica gel, reverse phase and SEPHADEX LH-20.

When the fermentation is carried out in a liquid medium, in one method, the mycelial solids are filtered and recovered from the fermentation medium. Alcohol is added to the mycelial cake, and the mycelial solid thoroughly mixed with the alcohol, filtered, and the filtrate collected and concentrated. In an alternative method, the whole broth can be extracted by the addition of one volume of alkanol, preferably methanol, and filtered to remove solid impurities. The alkanol extract is then adsorbed on DIAION SP-207 or other commercially available styrene-divinylbenzene copolymer. A second dilution/HP-20 adsorption/elution step is utilized to concentrate the sample in preparation for chromatographic separations. Sometimes, a third dilution/HP-20 adsorption/elution step may be desirable for volume reduction.

The alcoholic solvent to be employed in the initial extraction of the active agent from the solid nutrient medium or from the mycelial pad may be any of the lower alcohols such as methanol, ethanol, isopropanol, and the like. Methanol is preferred.

55 The water-immiscible non-polar organic solvent useful for extracting or partitioning the active agent from the alkanol or methanol solution are esters, such as ethyl acetate, isopropyl acetate, butyl acetate, or ketones, such as methyl ethyl ketone. Lower aliphatic esters are preferred.

The chromatographic separation may be carried out by employing conventional column chromatography with non-ionic resin or by high performance liquid chromatography employing reverse phase resin. The fractions containing the antibiotic Compound IA may be detected by bioautography using Candida albicans. Generally, more than one chromatographic separation steps are employed. In a most preferred procedure, one or more separations are carried out employing column chromatography and a final separation is carried out employing high performance liquid chromatography (HPLC) with C₁₈ reverse phase resin.

When conventional column chromatography is employed for chromatographic separations, silica gel is the preferred adsorbent. Usually more than one chromatographic separation is required. Silica gel may be used in

all the separations while employing different eluting agents. However, it may be combined advantageously with the use of a different adsorbent such as a dextran gel sold under the trade name of SEPHADEX LH-20 (Pharmacia). Other adsorbents such as alumina, styrene-divinylbenzene copolymers available commercially as DIAION HP-20, HP-30, HP-40, SP-207 (Mitsubishi Chemical Industries, Ltd.) and AMBERLITE XAD-2, XAD-4, XAD-16 (Rohm and Haas Co.) also may be employed.

In the fractionation and recovery of the active component by chromatography on silica gel, ester/alcohol mixtures with increasing concentration of alcohol provide good separations. A mixture of ethyl acetate and methanol has been found to be especially useful. These may be employed in isocratic, step gradient or continuous gradient systems. When a dextran adsorbent such as SEPHADEX LH-20, is employed, a chlorohydrocarbon/hydrocarbon/alcohol solvent system may be employed. A mixture of methylene chloride/hexane/methanol has been found to be especially useful.

In carrying out the HPLC separation, the alcohol solution containing material recovered from the conventional chromatography is concentrated and the residue dissolved in methanol and placed on a column packed with commercial reverse phase resin or on a column filled with silica gel/C₁₈ reverse phase resin prepared as amply reported in the literature. Alternatively, the alcohol solution may be diluted to 50 percent with water and pumped onto the column. The column is operated using methane/water (1:1 or optionally other ratios) at 800-2000 psi which produces a flow rate of about 20 ml/min. Separation is monitored at 210 nm.

The fractions are assayed for activity with *Candida albicans*. The product is recovered from the chromatographic procedures by combining the active fractions and concentrating under reduced pressure. The product then may be purified by conventional chromatography or preparative HPLC.

Other natural products which are embraced by the present invention include those which are described in the literature as echinocandins, aculeacins, mulundocandins, athlestatin (also referred to as an echinocandin T), sporiofungin or by number designations. The structures in some cases are not completely identified but the compounds are known to be neutral cyclohexapeptides with fatty acid side chain.

The echinocandins include echinocandin-A (also known as A-32204-A) produced by *Aspergillus nidulans*-*echinulatus* (Swiss patent 568,386); echinocandin-B (also known as A-32204-B and SF-7810-F) produced by *Asp. nidulans* (Swiss 568,368) and *Asp. rugulosus* (Helv. Chim. Acta 62 11252 (1979); German 2,549,127; Belg. 834,289); echinocandin-C (also known as SL-7810-FII) produced by *Asp. rugulosus* (Ger. 2,549,127; Belg. 834,289); echinocandin-D (also known as SL-7810-FIII) produced by *Asp. rugulosus* (Ger. 2,549,127; Belg. 834,289).

Echinocandins B, C and D may be produced together on the fermentations of a strain of *Aspergillus rugulosus*. The major metabolite in the fermentation of *Aspergillus rugulosus* is echinocandin B, having a molecular formula of C₅₂H₈₁N₇O₁₆. Echinocandin C and D which are the minor metabolites differ in certain of the amino acids making up the cyclic peptide ring. Thus, echinocandin C, having a molecular formula of C₅₂H₈₁N₇O₁₅, contains 3-hydroxyhomotyrosine in place of the 3,4-dihydroxyhomotyrosine present in echinocandin B. Echinocandin D, having a molecular formula of C₅₂H₈₁N₇O₁₃, contains 3-hydroxyhomotyrosine and ornithine instead of 3,4-dihydroxyhomotyrosine and 4,5-dihydroxyornithine.

Echinocandin B also may be produced by the cultivation of *Emmericella nidulans* as described in the working example.

The aculeacins include aculeacin-A, aculeacin-B, aculeacin-C, aculeacin-D, aculeacin-E, aculeacin-F, aculeacin-G, aculeacin-A A", aculeacin-DA", aculeacin-A G", aculeacin-A α and A γ , and aculeacin-D α and D γ . The aculeacins may be produced by the fermentation of *Aspergillus aculeatus* (either liquid or solid culture) (preferably M4214 FERM-P 2324) thereafter extracting the mycelium with a water miscible organic solvent concentrating and purifying by chromatography as described in U.S. Patent 3,978,210 and 4,212,858 German Patent 2,509,820 and Belgian Patent 826,393.

Another cyclohexapeptide antibiotic found in the literature is mulundocandin (R', R'', R''', R'''' are OH, R'''' is H, R is -CO(CH₂)₁₀-CH₃CH-CH₂CH₃) produced by *Asp. syndowic* No. Y-30462 (J. Antibiotics 40, 275 and 40, 281 (1987)). The antibiotic, which is present both in the culture filtrate and mycelium may be isolated from the ethyl acetate extract of a culture filtrate by silica gel column chromatography and droplet counter current chromatography as more fully described by Roy et al, J. Antibiotics 40, 275-280 (1987).

Still another echinocandin type cyclopeptide reported in the literature is sporiofungin (R', R'''' and R''' are OH, R'' is H, R'''' is CH₂CONH₂, R''' is H and R is a branched chain C₁₆ fatty acid radical,) which may be produced by cultivation of *Cryptosporiopsis* (WIPO publication, WO 82/00587) strain ATCC 20594 or NRRL 12192, on various conventional nutrient media as aerobic surface culture or immersion culture, and the mycelium separated from the culture broth and recovered in a conventional manner such as by homogenizing the mycelium with methanol, extracting from the methanol with a water-immiscible organic solvent such as ethyl acetate, removing the solvent and purifying by chromatography as more fully described in WO 82/00587.


Other echinocandin type cyclohexapeptides include X-73 produced by *Aspergillus rugulosus*, (Ind. J. B. Biochem. 7, 81 (1970)), athlestatin produced by *Aspergillus niger*, (Jap. J. Ant., 17 268 (1964), JP 66/12688; CA 65 19274; 70, 27607), S-31794-F-1 produced by *Acrophialophora limonispora* (U.S. 4,173,629; Ger. 2,628,965), A-30912-A or A-22082 produced by *Asp. nidulans* and *Asp. rugulosus* (U.S. 4,074,246; 4,074,245; 4,288,549; Ger. 2,643,485; CA 87, 20612) A-30912-B, A-30912-C, and A-30912-D, produced by *Asp. rugulosus* (U.S. 4,024,245).

Compounds which are modifications of natural products include those in which the side chain, i.e., R, when an unsaturated fatty acid, had been modified by reduction. This may be carried out catalytically, by using Pd on


carbon at atmospheric pressure.

Compounds in which R''' is -O-Alkyl (C₁-C₆) or -O-benzyl may be prepared by stirring together the corresponding compound in which R''' is OH with the appropriate alkanol under acidic conditions at ambient temperature for several hours or overnight to obtain the ether in the mixture and thereafter preferably quenching the reaction with dilute bicarbonate and recovering it by conventional procedures. In addition to the preparation in the working example, the preparation for a number of these compounds is more fully described in GB 2,050,385A.

Compounds in which R''' is



wherein R¹ is hydrogen, alkyl or substituted alkyl as previously defined, R² is hydrogen or C₁-C₄ alkyl, or R¹ and R² together form -(CH₂)₂-Q-(CH₂)₂- wherein Q is a connecting group such as -CH₂-, -NH- or -N(lower alkyl)- may be prepared by reacting a compound in which R''' is OH with a compound of the formula



in the presence of an organic acid such as p-toluenesulfonic acid, methanesulfonic acid or mineral acid in an aprotic solvent such as dimethylformamide as more fully described in Belgian 851,310.

Semi-synthetic compounds in which the acyl side chain, i.e., R in above formula, has been modified from naturally occurring fatty acids to structurally distinctive acyl groups may be produced by first removing the fatty acid side chain and thereafter introducing a distinctive acyl group. The fatty acid side chain preferably is removed enzymatically. The enzyme which is useful for deacylation of the cyclohexapeptides to obtain a cyclopeptide nucleus is that produced by certain microorganisms of the family Actinoplanaceae, especially the microorganism *Actinoplanes utahensis* NRRL 12052 which is available from the Northern Regional Research Center, USDA, Agricultural Research Service, Peoria Ill 61604, or as *A. utahensis* ATCC 14539 obtainable from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852.


The enzyme may be produced by growing *Actinoplanaceae* at temperatures between about 25°C and 30°C and a pH of between about 5.0 and 8.0, preferably between 6.0 and 7.0, with agitation and aeration for from about 40 to about 60 hours in a culture medium containing (a) an assimilable carbon source such as sucrose, glucose or glycerol, (b) a nitrogen source such as peptone, urea or ammonium sulfate (c) a phosphate source such as a soluble phosphate salt and (d) growth promoting inorganic salts.

In the deacylation, the cyclohexapeptide compound or substrate containing the cyclohexapeptide is added to the culture of *Actinoplanaceae* after the culture has been incubated for at least 48 hours. After addition of the substrate, the incubation of the culture is continued for about 24 hours or longer over temperatures in the range of from about 25°C to about 30°C.

The course of the reaction may be monitored by *Candida albicans* assay. The starting cyclohexapeptide compound is active against *C. albicans* but the deacylated nucleus compound is biologically inactive.

The deacylated nucleus compound then may be employed in the preparation of semi-synthetic compounds. Conventional acylation methods may be employed. In one preferred method, a 2,4,5-trichlorophenyl ester of the desired acid is reacted with the deacylated nucleus compound in an inert solvent such as dimethylformamide at room temperature for about 15 to 18 hours. This may be illustrated with a specific example as follows:

A deacylated nucleus of a compound the following structure wherein in formula (IB), E is H (Compound IBi)



may be prepared by first preparing a deacylating enzyme by the fermentation of *Actinoplanes utahensis* NRRL 12052 by first inoculating a slant with the organism, incubating the slant at 30° C for about 8 to 10 days and then using the slant growth to inoculate 50 milliliters of vegetative medium which is incubated at 30° C for about 72 hours with shaking. The incubated vegetative medium may be used to inoculate 400 milliliters of a second-stage vegetative medium having the same composition as the first stage vegetative medium and incubated at 30° C for about 48 hours with shaking. Thereafter, 800 milliliters of said medium may be used to inoculate 100 litres of production medium which is permitted to ferment at about 30° C for 42 hours with agitation and aeration to maintain dissolved oxygen level above 30 percent of air saturation at atmospheric pressure and to obtain therein the desired deacylating enzyme.

The medium for preparing the slant may be of the following composition: Baby oatmeal 60.0 g; yeast 2.5 g; K₂HPO₄ 1.0 g; Czepek's mineral stock 5.0 ml; agar 25.0 g; deionized water q.s. to 1 liter. Czepek's mineral stock is of the following composition: FeSO₄•7H₂O (in 2 ml. conc. HCl) 2 g; KCl 100 g; MgSO₄•H₂O 100 g; deionized water to 1 liter with final adjusted pH 6.7.

The medium suitable for both first stage and second stage vegetative medium may be of the following composition: Baby oatmeal 20.0 g; sucrose 20.0 g; yeast 2.5 g; dried grain 5.0 g; K₂HPO₄ 1.0 g; Czepek's mineral stock 5.0 ml and deionized water to 1 liter, final pH 6.8.

The production medium may be of the following composition: Peanut meal 10.0 g; soluble meat peptone 5.0 g; sucrose 20.0 g; KH₂PO₄ 0.5 g; K₂HPO₄ 1.2 g; MgSO₄•7H₂O 0.25 g; tap water q.s. to 1 liter.

An alcohol solution of a cyclohexapeptide (IB) where E is stearoyl (tetrahydroechinocandin B) or palmitoyl (aculeacin A) is added to the production fermentation medium above-described and the deacylation monitored by assay against *Candida albicans*.


On completion of the deacylation, the fermentation broth containing the nucleus compound is filtered, passed through HP-20 resin to place the nucleus compound on the column and the compound recovered by elution with water/methanol (7/3) at a rate of 200-300 ml/minute. The eluate is monitored for the nucleus compound by treating with acid chloride and assaying for activity against *Candida albicans*. The eluate fractions showing activity are concentrated to obtain the nucleus compound (IBI).

The deacylated cyclohexapeptide compound thus obtained is purified by dissolving in water/acetonitrile/acetic acid/pyridine (96/2/1/1) and purified by reversed phase liquid chromatography (LICHROPREP RP-18) at a flow rate of about 60 milliliters per minute and monitoring the separation at 280 nm using a UV monitor. On the basis of the absorption at 280 nm, the fractions containing the desired compound are combined, concentrated under reduced pressure and employed for acylation or lyophilized for subsequent use.

The preparation of the deacylated cyclohexapeptide compound having the same nucleus from echinocandin type natural products is more fully described in U.S. 4,293,482. Similar preparations of other deacylated cyclohexapeptide compounds with similar nuclei may be found described in U.S. patents 4,173,629; 4,293,490; 4,299,763; 4,299,762; and 4,304,716.

The deacylated compounds may then be employed to produce novel and/or unusual acylated compounds.

The acylation of the compound of the deacylated nucleus, to produce a unique acyl derivative may be illustrated with a preparation of a compound in which E in formula (IB) is



(Compound IBII). Such a compound would also be illustrated of a compound of formula (I) in which R is a radical defined under (C)(i).

In the acylation, the acyl group is preferably introduced by means of a trichlorophenyl ester. Thus, first, the trichlorophenyl ester may be prepared by treating the side chain acid with 2,4,5-trichlorophenol in the presence of a coupling agent such as N,N'-dicyclohexylcarbodiimide in an inert solvent such as methylene chloride. About 1.1 mole of the 2,4,5-trichlorophenol and 1.0 mole of the coupling agent is used for each mole of the alkoxybenzoic acid. The mixture is stirred at room temperature for 15 to 18 hours to obtain the ester which may be recovered by filtering the solid and evaporating the filtrate under reduced pressure, then recrystallizing the residue.

The ester thus prepared is added to a solution of the nucleus compound in dimethylformamide and stirred for about 18 hours and then the solvent evaporated off. The residue is washed and then chromatographed on silica gel using ethyl acetate-methanol (3/2) as eluant to obtain the desired octyloxybenzoyl derivative which may be represented in formula IB with the octyloxybenzoyl group as E. Such a compound may be named 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithine]echinocandin B (Compound IBII).

In addition to Compound IA, other preferred compounds include Compound IBII; echinocandin B (R', R'', R''', R'''' are OH, R'''' and R''' are CH₃; R is

$\text{-}\overset{\text{O}}{\parallel}\text{C}(\text{CH}_2)_4\text{CH}=\text{CH}(\text{CH}_2)_7\text{CH}_3$; tetrahydroechinocandin B (R', R'', R''', R'''' are OH; R'''' and R''' are CH₃; R is

$\text{-}\overset{\text{O}}{\parallel}\text{C}(\text{CH}_2)_{16}\text{CH}_3$); O-methyltetrahydroechinocandin B (R', R'', and R''' are OH; R'' is -OCH₃; R'''' and R''' are CH₃; R is

$\text{-}\overset{\text{O}}{\parallel}\text{C}(\text{CH}_2)_{16}\text{CH}_3$); and O-benzyltetrahydroechinocandin B (R', R'' and R''' are OH, R'' is -OCH₂C₆H₅, R'''' and R''' are CH₃; R is

$\text{-}\overset{\text{O}}{\parallel}\text{C}(\text{CH}_2)_{16}\text{CH}_3$).

The method of the present invention for the treatment of or for the prevention of *Pneumocystis carinii* infections in mammals comprises administering to subjects infected with or immune compromised subjects susceptible to being infected with *Pneumocystis carinii*, a therapeutically effective or anti-infective amount of a cyclohexapeptide compound represented by Formula I.

The efficacy of the cyclohexapeptide compounds for therapeutic and anti-infective purposes in immune-compromised patients may be determined first in studies on immunosuppressed rats and immunosuppressed mice.

In a representative study employing Compound IA in rats, nine male Sprague-Dawley rats weighing approximately 200 grams each, were immunosuppressed by the addition of dexamethasone to the drinking water (2.0 mg/liter) for six weeks to induce the development of *P. carinii* infections. To enhance the infection the rats were also maintained on a low protein diet. At the beginning of the seventh week the rats were divided into three groups of three rats each. All three groups continued to receive dexamethasone in the drinking water and low protein diet throughout the remainder of the study. The rats in Group I were kept as untreated infected controls, those in Group II were injected intraperitoneally twice daily for two weeks with 0.5 ml of sterile dH₂O containing 2 mg of Compound IA, and those in Group III were treated with trimethoprim-sulfamethoxazole (TMP-SMZ) in the drinking water (208 mg TMP and 1.040 g of SMZ/liter) for two weeks, a known treatment for *P. carinii* infections. Two rats died early in the experiment. One was from group II (rat 72A) and the other from group III (rat 75A). It was not determined whether they died from *Pneumocystis pneumonia*.

At the end of the two week treatment period (a total of eight weeks of immunosuppression) the animals were sacrificed and the lung tissue removed. The lungs from each animal were weighed, and then processed to determine the number of cysts and parasite nuclei for each animal. The results of this experiment are shown in Table I.

TABLE I
IN VIVO TESTING OF COMPOUND IA & TMP-SMZ FOR ACTIVITY AGAINST P. CARINII

RAT #	TOTAL CYSTS		TOTAL NUCLEI		LUNG WT. (gms)	CYST SCORE
	per lung	per gm lung	per lung	per gm lung		
GROUP I (control)						
714	3.2x10 ⁸	1.0x10 ⁸	5.8x10 ⁹	1.8x10 ⁹	3.16	4
74A	1.5x10 ⁹	4.5x10 ⁸	1.4x10 ¹⁰	4.2x10 ⁹	3.32	4+
74B	7.5x10 ⁸	2.1x10 ⁸	1.4x10 ¹⁰	3.9x10 ⁹	3.56	4+
GROUP II (COMPOUND IA)						
71B	N.D.*	N.D.*	2.1x10 ⁹	7.4x10 ⁸	2.83	0
72B	N.D.*	N.D.*	1.0x10 ⁹	7.1x10 ⁸	1.40	0
73B	6.0x10 ⁸	5.4x10 ⁶	2.0x10 ⁸	1.8x10 ⁸	1.11	1
75B	5.4x10 ⁸	2.3x10 ⁶	1.6x10 ⁹	7.0x10 ⁸	2.28	1

*N.D. = No cysts were detected in 50 fields, indicating that there were less than 2×10^5 (limits of resolution).

These results demonstrate the Compound IA has efficacy against *P. carinii* infections. No cysts could be found in the lungs of either rat treated with Compound IA as compared to levels ranging from 3.2×10^8 - 1.5×10^9 in the control animals. There was also a reduction in the number of parasite nuclei (1.0 - 2.1×10^9) in these animals as compared to controls (5.8×10^9 - 1.4×10^{10}). These results are superior to those seen in animals treated with TMP-SMZ, a known treatment for *P. carinii* infections, since the TMP-SMZ rats had detectable levels of *P. carinii* cysts.

A similar rat study was conducted to determine the antipneumocystis activity of 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ii ornithine]echinocandin B (Compound IBII). After 6 weeks of immunosuppression the rats were injected twice daily for 7 days with vehicle alone. Compound IBII (2 mg/kg) or Compound IA (2 mg/kg) during which time immunosuppression was continued. At the end of the treatment period the animals were sacrificed and the number of cysts per lung determined. Compound IA reduced the number of cysts by > 99% while Compound IBII reduced the cyst counts by approximately 84% (compared to the vehicle control, 10% DMSO).

Studies were also carried out with mice and still other compounds within Formula I. Representatives of the method used in the studies carried out with mice and with other compounds is as follows:

Male C3H/HeN mice, weighing 22-24 grams each, were immunosuppressed by the addition of dexamethasone to the drinking water for six weeks to induce the development of *P. carinii* infections. To enhance the infections the mice were also maintained on a low protein diet. At the beginning of the seventh week, the mice were randomly divided into groups of 6. All groups continued to receive dexamethasone in the drinking water and low protein diet for the remainder of the study. Mice in each group were injected intraperitoneally twice daily with 0.5 ml of a 10% DMSO solution as a vehicle control or the same cocktail containing drug to achieve a dose of 2 mg/kg. This procedure was continued for one week.

At the end of the treatment period (a total of seven weeks of immunosuppression) the animals were sacrificed and the lung tissue removed. The tissue was then processed to determine the number of cysts for each animal.

The results for the various compounds are seen in Table II.

TABLE II

Compound	Dosage	% Reduction of Cysts
Tetrahydroechinocandin B (TEB)	2mg/kg	99%
O-Methyl-TEB	2mg/kg	> 99%
O-Benzyl-TEB	2mg/kg	99%
Echinocandin B	2mg/kg	> 99%
Compound IA	2mg/kg	> 99%

From the foregoing test results and from known dosage ranges for TMP-SMZ as applied to man, it is determined that generally from about 0.5 to about 20.0 mg/kg of body weight of the cyclohexapeptide anti-pneumocystis agent (Compound I) may be employed while considering the patient's health, weight, age and other factors which influence response to a drug as well as whether it is to be applied to a human patient or to an animal. These amounts, when expressed as doses suitable for human beings, are in the range of from about 35 mg to about 1500 mg daily preferably by parenteral administration.

The outstanding properties are most effectively utilized when the compound is formulated into novel pharmaceutical compositions with a pharmaceutically acceptable carrier according to conventional pharmaceutical compounding techniques.

The novel compositions contain at least 1% by weight of the active compound. In preparing the compositions, Compound I is intimately admixed with any of the usual pharmaceutically acceptable carrier.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), pulmonary (nasal or buccal inhalation), nasal administration, or insufflation.

The cyclohexapeptide anti-pneumocystis agent, Compound I, is preferably formulated into compositions for injection and may be presented in unit dosage form in ampoules or in multidose containers, if necessary with an added preservative. The compositions may also take such forms as suspensions, solutions or emulsions in oil or in vehicles such as 20/80 ethanol/propylene glycol or 20-35% polyethylene glycol 300 and may contain formulating agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredients may be in powder form for reconstituting with a suitable vehicle prior to administration or for use in insufflation.

The term "unit dosage form" as used in the specification and claims refer to physically discrete units, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the pharmaceutical carrier. Examples of such unit dosage forms measured units in ampoules or in multidose containers and the like. A unit dosage of the present invention will generally contain from 100 to 1000 milligrams of the cyclohexapeptide anti-pneumocystis agent.

Any suitable route of administration may be employed for providing a patient with an effective dosage of a

compound of Formula I. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, disperions, suspensions, solutions, capsules, creams, ointments, aerosols, powders for insufflations and the like.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), pulmonary (nasal or buccal inhalation), nasal administration, or insufflation.

For administration by inhalation, the compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The compounds may also be delivered as powders which may be formulated and the powder compositions may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery system for inhalation is a metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of Compound I in suitable propellants, such as fluorocarbons or hydrocarbons.

Because of the difficult solubility of the compound in pharmaceutically acceptable liquid carriers and because of desirability to directly treat lung and bronchi, aerosol administration is a preferred method of administration. Insufflation is also a desirable method, especially where infection may have spread to ears and other body cavities.

Alternatively, parenteral administration may be employed using drip intravenous administration. In such method, the drug may be solubilized in alcohol/propylene glycol or polyethylene glycol compositions. Intraperitoneal injection also may be employed.

The following examples illustrate the preparation of Compound IA and other lipophilic cyclohexapeptide compound of Formula I and the compositions useful in the present invention of controlling Pneumocystis carinii but are not to be construed as limiting.

PREPARATION OF COMPOUND IA

EXAMPLE A

Fermentation

A frozen culture in glycerol of Isolate 2 of Culture 8525-307P originally isolated from water, identifiable as Zalerion Arboricola, ATCC 20868, and maintained in the Merck culture collection was employed in the fermentation.

A 2 milliliter portion of the frozen culture was defrosted and aseptically transferred to a 250 milliliter unbaffled Erlenmeyer flask containing 54 milliliters of Medium 1. Medium 1, after inoculation, was incubated at 28°C with rotary agitation (220 rpm, 2" throw shaker) for three days. At the end of this period, 2.0 milliliters of the growth medium were aseptically transferred to each of several unbaffled 250 milliliter Erlenmeyer flasks containing Medium 1. The inoculated flasks were incubated at 28°C for 2 days.

12.5 milliliters of the mature seed broth were inoculated into five production flasks containing Medium 2, and incubated at 25°C for seven days under static conditions to obtain an antibiotic compound on the fermentation medium.

The media employed in the foregoing fermentation were:

EP 0 359 529 A1

MEDIUM 1 (KF Seed Medium)

	Corn Steep Liquor	5 g
	Tomato Paste	40 g
5	Oat Flour	10 g
	Glucose	10 g
	Trace Elements Mix	10 ml
	Distilled Water	to 1000 ml
	pH 6.8	
10	Trace Elements Mix:	
	FeSO ₄ ·7H ₂ O	1 g
	MnSO ₄ ·4H ₂ O	1 g
	CuCl ₂ ·2H ₂ O	25 mg
15	CaCl ₂	100 mg
	H ₃ BO ₃	56 mg
	(NH ₄) ₆ MoO ₂ ·4H ₂ O	19 mg
	ZnSO ₄ ·7H ₂ O	200 mg
20	Distilled Water	to 1000 ml

MEDIUM 2 (F204 Solid Medium)

25		Amount/flask
	Millet Base	15 g
	Yeast Extract	.5 g
	Sodium tartrate	.1 g
30	Ferrous Sulfate Crystals	.01 g
	Monosodium Glutamic Acid	.1 g
	Corn Oil	.1 ml

35

Isolation

40 Five hundred milliliters of methanol were added to each of the five 2-liter flasks of solid phase fermentation. The contents of the flask were then combined and stirred to extract methanol-soluble material and the mixture then filtered. The spent cake was stirred with an additional 2500 milliliters of methanol to further extract methanol soluble material and the mixture then filtered.

The filtrate and wash were combined and concentrated to 500 milliliters.

45 The aqueous methanolic concentrate thus obtained then was extracted with two 500 milliliter portions of ethyl acetate. The spent aqueous solution was placed onto a DIAION HP-20 column to adsorb the active material thereon and the latter then eluted therefrom with methanol. The eluates were combined with the previously obtained ethyl acetate extracts and the combined ethyl acetate solutions were concentrated to dryness. The residue was chromatographed on 200 milliliters of SEPHADEX LH-20 using 5:5:2 methylene chloride/hexane/methanol as eluant.

50 The fractions active as determined by *Candida albicans* were combined and chromatographed on 200 milliliters of silica gel (EM Science, KEISELGEL 60, 230-400 mesh) using a step gradient elution with ethyl acetate/methanol. The active fractions from this chromatography were combined, concentrated and chromatographed on silica gel using a 75:25 ethyl acetate/methanol isocratic system.

55 The active portions from this chromatography were then combined and placed on 100 milliliters of SEPHADEX LH-20 using methanol as eluting solvent. The eluate, after vaporization of the solvent, yielded 95 milligrams of a purified compound. The compound was a white solid having a ¹HNMR spectrum as seen in Figure 1.

60

EXAMPLE B

65

Fermentation

In a manner similar to that described in Example A, the contents of one frozen vial of ATCC 20868 from the Merck culture collection were defrosted and aseptically transferred to a 250 milliliter unbaffled flask containing 54 milliliters of KF medium (Medium 1) containing 0.4 percent agar. The modified Medium 1, after inoculation, was incubated at 28°C with 220 rpm agitation for 48 hours. At the end of this period, 10 milliliters of the growth medium was transferred to a 2-liter unbaffled flask containing 500 milliliters of KF medium containing 0.4 percent agar. After inoculation, the resulting medium was incubated for 24 hours at 28°C with 220 rpm agitation.

Twenty 2-liter flasks each containing 120 grams of Medium 2 and 120 milliliters of a stock solution consisting of

Yeast extract	5 parts by weight
Sodium tartrate	1 part by weight
Ferrous sulfate crystals	0.1 part by weight
Monosodium glutamic acid	1 part by weight
Corn oil	1 part by weight

were autoclaved for 20 minutes at 122°C and then reautoclaved with 80 milliliters of water for another 20 minutes at 122°C. The flasks were allowed to cool, then inoculated with 20 milliliters of seed medium prepared as above described, and the inoculated flasks incubated at 25°C under static conditions for 14 days.

Isolation

To each of nineteen 2-liter solid fermentation flasks was added 1 liter of methanol and the contents combined, stirred and filtered. The spent cake was extracted twice with 6 liters of methanol. The aqueous methanol filtrates were then concentrated and the concentrate extracted twice with 3 liters of ethyl acetate. The ethyl acetate extracts were combined, dried and concentrated to about 100 milliliters.

The concentrate was then coated on silica gel by adding 100 milliliters of methanol and 100 milliliters of silica gel thereto, intimately contacting the components and then removing the solvent on a rotary evaporator. The dried silica gel was then applied to a column of 500 milliliters of silica gel, the column washed with ethyl acetate to remove impurities and eluted with 9:1 ethyl acetate/methanol. The eluates containing antibiotic material testing positive against *Candida albicans* were recovered and combined.

The antibiotic rich cut from the silica gel chromatography was dissolved in 200 milliliters of 10:10:1 methylene chloride/hexane/methanol and the resulting solution combined with 40 milliliters of SEPHADEX LH-20 (which previously had been prepared by soaking overnight in methanol followed by washing twice with 200 milliliters of methylene chloride/hexane/methanol). After a few minutes, the supernatant was removed by filtration and the SEPHADEX LH-20 was washed with 200 milliliters of methylene chloride/hexane/methanol and then filtered. The filtrates were found not to contain the active constituent, and were discarded. The active constituent which had partitioned into the dextran SEPHADEX LH-20 beads was extracted therefrom by washing twice with 200 milliliters of methanol, and these methanol washes were combined and concentrated.

The methanol concentrate was next applied to 200 milliliters of silica gel and eluted with 75:25 ethyl acetate/methanol and the eluates combined and the solvent vaporized to obtain Compound 1A. Compound 1A is a white powder having a decomposition point of 206-214°C.

PREPARATION OF ECHINOCANDIN B DERIVATIVES

EXAMPLE C

Preparation of Echinocandin B

A. Fermentation

Culture MF 5100 *Emericella nidulans* ATCC 20956 from the Merck culture collection was used for seed train development as follows: One frozen vial (about 2.5 ml) of MF 5100 was used to inoculate 54 mL of KF medium and cultured in 250 ml unbaffled Erlenmeyer flasks at 28°C and 220 rpm for 72 hours to obtain "B" stage culture. Five mL of "B" stage culture was used to inoculate 500 mL of KF medium in 2 liter unbaffled Erlenmeyer flask and incubated at 28°C and 220 rpm for 48 hours to obtain "C" stage cultures. Four "C" stage flasks were used to inoculate 160 liters of KF medium with trace element mix #2 in a "D" stage (300 liter stirred fermenter).

EP 0 359 529 A1

The three stages of seed train were used for "E" scale fermentation in an 800 liter fermenter. The fermentation was carried out for 23 hours at 28°C, back pressure 0.7 kg/cm², aeration with 0.300m (53 liters per minute) and agitation of 100 rpm.

The production "E" stage was inoculated with 5% seed from "D" stage in 5700 liters of a production medium of the following composition:

	Glycerol	85 g/L
	Pectin	10 g/L
	Peanut Meal	4 g/L
10	Peptonized Meal	4 g/L
	Peptonized Milk	4 g/L
	Tomato Paste	4 g/L
	Corn Steep Liquor	4 g/L
15	Lard Water	4 g/L
	Glycine	2 g/L
	Potassium phosphate monobasic	2 g/L
20	Polyglycol P-2000	1 mL/L

The medium was sterilized for 25 minutes at 123°C and the fermentation then carried out at temperature of 28°C, positive pressure of about 0.8 kg/cm², aeration 6000 liters per minute, agitation speed of 120 rpm; pH in the range 5.5-6.5 with dissolved oxygen maintained above 30 percent for about 36 hours.

25 B. Isolation

The whole broth from the above fermentation containing 355 grams of echinocandin B by HPLC assay (against reference standard echinocandin B) was extracted overnight with an equal volume of methanol. The diluted broth was centrifuged to remove broth solids and the centrifugate adsorbed on a 450 liter SP 207 column which had been equilibrated with 50:50 methanol/water. After a 1000 liter water wash and a 1000 liter 50:50 methanol/water wash, the column was eluted with methanol and the eluate concentrated to 189 liters and partitioned with methylene chloride. The aqueous methanol layer was adjusted to 50:50 methanol/water by the addition of water and adsorbed on a 100 liter HP-20 column prewashed with methanol and pre-equilibrated with 50:50 methanol/water. After a 50:50 methanol/water wash, the column was eluted with 80:20 methanol/water. The 80:20 eluate was adjusted to 50:50 with water, and then was adsorbed on a 27 liter SP207 column, preequilibrated as before. The column was then eluted with methanol.

A 19 g aliquot of SP 207 eluate was diluted to 50:50 methanol/water with water and adsorbed on a 3.9 liter Amicon preparative C-18 column. A gradient of acetonitrile/water (25:75) to 100 percent acetonitrile over a 75 minute period was used to elute the compound. Selected fraction after concentration and lyophilization from acetonitrile/water provided 16 grams of Echinocandin B of 87-91 percent purity.

The identity and purity of the foregoing echinocandin B was determined by HPLC and spectral comparison with a sample of Echinocandin B, of the properties below, previously prepared and purified, and its identity established by comparison with a published structure of Echinocandin B.

45 EXAMPLE D

50 Preparation of Tetrahydroechinocandin B

5.0 Grams of Echinocandin B (EB) of about 87 percent purity (HPLC) was dissolved in 150 milliliters of absolute ethanol and was added to 10% Pd-C in 80 milliliters of ethanol which had been prerduced under 1 atmosphere of hydrogen for 20 minutes. After the addition, the mixture was stirred rapidly under 1 atmosphere of hydrogen for 4 hours. Analysis by HPLC using CH₃CN/H₂O (81/19), on a C8 ZORBAX column at flow rate 1 ml/min at 30°C showed that the reaction was virtually complete at this time.

The reduction mixture was filtered through a Celite pad, the pad washed with ethanol and the mixture was concentrated in vacuo to obtain a solid. The solid was purified in ten 0.5 gram batches dissolved in 5 milliliters of CH₃CN/H₂O (81/19). The HPLC was carried out on C8 ZORBAX 1 inch column and eluted with 80% methanol/20% water at a flow rate of 15 ml/min. A total yield of 3.15 grams (79%) of tetrahydroechinocandin B was obtained.

65 EXAMPLE E

Preparation of O-Methyltetrahydroechinocandin B

55 Milligrams of tetrahydroechinocandin B was dissolved in 4 milliliters of methanol. To the solution was added 4.0 milligrams of p-toluenesulfonic acid and the mixture stirred at room temperature for 3 hours. HPLC analysis indicated that the reaction was complete at this time.

The reaction mixture was quenched with 1N sodium bicarbonate solution and washed twice with saturated sodium chloride solution and then diluted with 200 milliliters of isopropanol/chloroform (2/7).

The organic layers were dried over sodium sulfate, filtered and the solvent vaporized to obtain a near white solid as residue. The latter was purified first with methanol, and then with H₂O/CH₃CN (2:3) and chromatographing by MPLC in H₂O/CH₃CN (2:3) on a C8 LOBAR size A column in 4 milliliter fractions. The compound was obtained in a yield of 42.4 mg. (76%)

EXAMPLE F

Preparation of O-Benzyltetrahydroechinocandin B

50 Milligrams of tetrahydroechinocandin B was suspended in 1 milliliter of tetrahydrofuran. To it was added 97 milliliters of anhydrous benzyl alcohol and the resulting mixture stirred until all was in complete solution. To the resulting solution was added 1 mg. of p-toluenesulfonic acid monohydrate and the mixture then stirred at room temperature for 3 hours whereupon a HPLC analysis showed that the reaction was complete. The reaction mixture was diluted with 1N sodium bicarbonate solution and then with isopropyl alcohol/chloroform (3:7). The organic solution was washed twice with water, dried and evaporated to recover the product as residue.

The following examples illustrate novel compositions useful in the practice of the present invention but are not to be construed as limiting.

Aerosol compositions may be prepared having the following formulations:

EXAMPLE I

	<u>Per Canister</u>
Compound Ia	24 mg
Lecithin, NF Liquid Concentrate	1.2 mg
Trichlorofluoromethane	4.025 g
Dichlorodifluoromethane	12.15 g

EXAMPLE II

	<u>Per Canister</u>
Echinocandin B	24 mg
Oleic acid	1.2 mg
Trichlorofluoromethane	4.025 g
Dichlorodifluoromethane	12.15 g

EXAMPLE III

	<u>Per Canister</u>
O-Methyltetrahydroechinocandin B	24 mg
Sorbitan trioleate	1.2 mg
Trichlorofluoromethane	4.025 g
Dichlorodifluoromethane	12.15 g

An injectable suspension (1M) may be prepared as follows:

EXAMPLE IV

	mg/ml
Compound Ia	10
5 Methylcellulose	5.0
Tween 80	0.5
Benzyl alcohol	9.0
Benzalkonium chloride	1.0
10 Water to 1 ml.	

General Information

For Echinocandin B

20 *Emericella nidulans* (a later growth stage of *Aspergillus nidulans*, established as a producing organism for Echinocandin B) and maintained as MF 5100 in the Merck culture collection was employed in the fermentation. A culture was deposited with the American Type Culture Collection under the Budapest Treaty and assigned ATCC No. 20956.

25 Echinocandin B, prepared as described in Example C was compared for percent purity against previously prepared Echinocandin B having the following MS and NMR spectra and identified by comparison with published structure for Echinocandin B.

Mass Spectral Data: FAB mass spectra were recorded on a VG 20-253 mass spectrometer. A matrix of dithiothreitol-dithiocrythritol-LII was used and M+Li was observed at m/z 1066 corresponding to the molecular weight 1059.


30 ¹H NMR Spectrum: The spectrum, Fig. 2, was recorded in CD₃OD on a Varian XL-300 NMR Spectrometer at 300 MHz. Chemical shifts are shown in ppm relative to tetramethylsilane (TMS) at zero ppm using the solvent peak at 3.30 ppm as internal reference.

35 ¹³C NMR Data: The spectrum was recorded in CD₃OD at 24°C on a Varian XL-400 NMR spectrometer at 100 MHz. Chemical shifts are given in ppm relative to tetramethylsilane at zero ppm using the solvent peak at 49.0 ppm as internal reference.

¹³C Chemical Shifts (CD₃OD, 100 MHz): 11.28, 14.43, 19.66, 20.04, 23.62, 26.54, 27.05, 28.15, 28.20, 30.27(x2), 30.42, 30.46, 30.78, 32.65, 35.08, 36.82, 38.60, 39.06, 51.67, 52.94, 56.32, 56.84, 57.19, 58.61, 62.46, 68.33, 69.59, 69.70, 70.87, 71.29, 74.60, 75.52, 75.77, 76.94, 116.21(x2), 129.06(x2), 129.60(x2), 130.94, 130.96, 133.09, 158.46, 169.97, 172.48, 172.51, 172.79, 173.53, 174.35, 176.19.


Claims

45 1. The use of a cyclohexapeptide compound represented by the formula



65 wherein
R' is H or OH;

R'' is H or OH;
R''' is H, OH, -O-alkyl(C₁-C₆), -O-benzyl, or



wherein R¹ is H, C₁-C₁₂ alkyl, C₃-C₇ cycloalkyl, hydroxyethyl, lower alkoxyethyl, furylmethyl, tetrahydrofurylmethyl, phenyl, phenylalkyl wherein the phenyl may be substituted with halo, hydroxy, lower alkyl and lower alkoxy; R² is hydrogen or lower alkyl; or R¹ and R² together form -(CH₂)₂Q(CH₂)₂- wherein Q is a connecting group such as -CH₂-, -NH- or -N- lower alkyl; and m is from 0 to 4;

R''' is H or OH;



R'''' is H, CH₃ or CH₂CONH₂;

R'''' is H or CH₃; and


R is (A) - $\text{C}(=\text{O})$ -alkyl from 6 to 24 carbon atoms

(B) - $\text{C}(=\text{O})$ -alkenyl from 6 to 24 carbon atom

(C) wherein R', R'', R''' and R'''' are OH, R'''' is CH₃ then



or



wherein A is divalent oxygen, sulfur, sulfinyl or sulfonyl;

A' is divalent oxygen, sulfur, sulfinyl, sulfonyl or -NH-;

X is hydrogen, chloro, bromo, iodo, nitro, C₁-C₃ alkyl, hydroxy, C₁-C₃ alkoxy, mercapto, C₁-C₃ alkylthio, carbamyl or C₁-C₃ alkylcarbamyl;

X' is chloro, bromo or iodo,

R₁ is hydrogen, C₁-C₁₈ alkyl or C₂-C₁₈ alkenyl;

W is C₁-C₁₀ alkylene or C₂-C₁₀ alkenylene;


m, n and p are 0 or 1, but if m is 0, n must be 0;

provided that the sum of the carbon atoms in R₁ and W groups must be greater than 4 but cannot exceed 21; that when X is mercapto, A and A' cannot be sulfinyl or sulfonyl; and that when A and A' are sulfinyl or sulfonyl, they must be in equal oxidation states;


(D) when R' is OH, R'', H, then



wherein Y is a divalent aminoacyl radical of the formula




wherein Z is C₁-C₁₀ alkylene or C₅-C₈ cycloalkylene



wherein R₃ is hydroxymethyl, hydroxyethyl, mercaptomethyl, mercaptoethyl, methylthioethyl, 2-thienyl,

3-indolomethyl, phenyl, benzyl, halophenyl, halobenzyl, C₁-C₃ alkylphenyl, C₁-C₃ alkylbenzyl, C₁-C₃ alkylthiophenyl, C₁-C₃ alkylthiobenzyl, carbamylphenyl, carbamylbenzyl, C₁-C₃ alkylcarbamylphenyl or C₁-C₃ alkylcarbamylbenzyl;

(c)




wherein X² is hydrogen or halo; for the preparation of a medicament, useful for the treatment of or for the prevention of *Pneumocystis carinii* infections.

2. The use according to Claim 1 wherein the cyclohexapeptide compound is administered at a dose of from 0.5 to 20.0 mg/kg of mammalian body weight.

3. The use according to Claim 1 wherein the cyclohexapeptide compound is administered to a patient with an impaired immune system infected with *Pneumocystis carinii*.

4. The use according to Claim 1 wherein the cyclohexapeptide compound is administered to a patient infected with *Pneumocystis carinii* and exhibits AIDS.

5. A composition useful for the treatment of *Pneumocystis carinii* infections in mammals comprising a pharmaceutically acceptable carrier and a cyclohexapeptide compound having the formula




wherein

R' is H or OH;

R'' is H or OH;

R''' is H, OH, -O-alkyl(C₁-C₆), -O-benzyl, or



wherein R¹ is H, C₁-C₁₂ alkyl, C₃-C₇ cycloalkyl, hydroxyethyl, lower alkoxyethyl, furylmethyl, tetrahydrofurylmethyl, phenyl, phenylalkyl wherein the phenyl may be substituted with halo, hydroxy, lower alkyl and lower alkoxy; R² is hydrogen or lower alkyl; or R¹ and R² together form -(CH₂)₂Q(CH₂)₂- wherein Q is a connecting group such as -CH₂-, -NH- or -N(lower alkyl)-; and m is from 0 to 4;

R''' is H or OH;


R'''' is H, CH₃ or CH₂CONH₂; and

R'''' is H or CH₃; and


R is (A) -C(=O)-alkyl from 6 to 24 carbon atoms

(B) -C(=O)-alkenyl from 6 to 24 carbon atom

(C) when R', R'', R''' and R'''' are OH, R'''' is CH₃ then




5



or

10



15

20

wherein A is divalent oxygen, sulfur, sulfinyl or sulfonyl;
 A' is divalent oxygen, sulfur, sulfinyl, sulfonyl or -NH-;
 X is hydrogen, chloro, bromo, iodo, nitro, C₁-C₃ alkyl, hydroxy, C₁-C₃ alkoxy, mercapto, C₁-C₃ alkylthio, carbamyl or C₁-C₃ alkylcarbamyl;
 X' is chloro, bromo or iodo,
 R₁ is hydrogen, C₁-C₁₈ alkyl or C₂-C₁₈ alkenyl;
 W is C₁-C₁₀ alkylene;
 m, n and p are 0 or 1, but if m is 0, n must be 0;
 provided that the sum of the carbon atoms in the R₁ and W groups must be greater than 4 but cannot exceed 21; that when X is mercapto A and A' cannot be sulfinyl or sulfonyl; and that when A and A' are sulfinyl or sulfonyl, they must be in equal oxidation states;
 (D) when R' is OH, R'', H, then


25

30




35

wherein N is a divalent aminoacyl radical of the formula



wherein Z is C₁-C₁₀ alkylene or C₅-C₆ cycloalkylene

40




45

wherein R₃ is hydroxymethyl, hydroxyethyl, mercaptomethyl, mercaptoethyl, methylthioethyl, 2-thienyl, 3-indolomethyl, phenyl, benzyl, halophenyl, halobenzyl, C₁-C₃ alkylphenyl, C₁-C₃ alkylbenzyl, C₁-C₃ alkylthiophenyl, C₁-C₃ alkylthiobenzyl, carbamylphenyl, carbamylbenzyl, C₁-C₃ alkylcarbamylphenyl or C₁-C₃ alkylcarbamylbenzyl;

(c)

50



55

wherein X² is hydrogen or halo.

6. A composition according to Claim 5 which is suitable for administration by inhalation.


7. A composition according to Claim 5 which is suitable for parenteral administration.

60


8. A composition according to Claim 5 in unit dosage form.


9. The use according to Claim 1 wherein the compound is that having the formula

65



- 25
10. The use according to Claim 1 wherein the compound is 1-[(4R,5R)-4,5-dihydroxy-N²-(10,12-dimethyl-1-oxotetradecyl)-L-ornithine-5-(threo-3-hydroxy-1-glutamine)echinocandin B.
11. The use according to Claim 1 wherein the compound is 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithine]echinocandin B.
12. The use according to Claim 1 wherein the compound is echinocandin B.
13. The use according to Claim 1 wherein the compound is tetrahydroechinocandin B.
- 30
14. The use according to Claim 1 wherein the compound is O-methyltetrahydroechinocandin B.
15. The use according to Claim 1 wherein the compound is O-benzyltetrahydroechinocandin B.
- 35
- 40
- 45
- 50
- 55
- 60
- 65







European Patent
Office

EUROPEAN SEARCH REPORT

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 89309257.7
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D, P, A	<u>EP - A2 - 0 311 193</u> (MERCK) * Claims 1, 3, 4, 5 * --	1, 5, 9, 10, 12	A 61 K 37/02 C 07 K 7/64
D, P, A	<u>EP - A2 - 0 311 194</u> (MERCK) * Claim 1 * --	1, 9, 10	
A	<u>DE - A1 - 2 803 581</u> (SANDOZ) * Claims 1, 3 * --	1, 5, 13	
A	<u>EP - A1 - 0 032 009</u> (ELI LILLY) * Claim 1; page 1, lines 1-28 * -----	1	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A 61 K 37/00 C 07 K 7/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 29-12-1989	Examiner DUNGLER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



(11) Publication number : **0 359 529 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification :
03.03.93 Bulletin 93/09

(51) Int. Cl.⁵ : **A61K 37/02, C07K 7/64**

(21) Application number : **89309257.7**

(22) Date of filing : **12.09.89**

(54) **Method for the control of pneumocystis carinii.**

(30) Priority : **12.09.88 US 242767**
16.09.88 US 246042
30.08.89 US 398889

(43) Date of publication of application :
21.03.90 Bulletin 90/12

(45) Publication of the grant of the patent :
03.03.93 Bulletin 93/09

(84) Designated Contracting States :
AT BE CH DE ES FR GB IT LI LU NL SE

(56) References cited :
EP-A- 0 032 009
EP-A- 0 311 193
EP-A- 0 311 194
DE-A- 2 803 581

(73) Proprietor : **MERCK & CO. INC.**
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900 (US)

(72) Inventor : **Schmatz, Dennis M.**
306 Elizabeth Avenue
Cranford New Jersey 07016 (US)

(74) Representative : **Hesketh, Alan, Dr. et al**
European Patent Department Merck & Co.,
Inc. Terlings Park Eastwick Road
Harlow Essex, CM20 2QR (GB)

EP 0 359 529 B1

Note : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

BACKGROUND OF THE INVENTION

5 Pneumocystis carinii is an opportunistic organism which is widely prevalent but generally dormant until bodily defenses of the host are compromised whereupon it propagates in the host causing disease. Generally it is present in the lungs although extrapulmonary infections have been reported. Lung infections with this organism in immunocompromised individuals generally lead to pneumonia and are usually fatal if untreated. It has been suggested that patient to patient transmission also may occur in immunocompromised individuals.

10 The immunocompromised state of individuals is usually associated with genetic defects, lymphoproliferative diseases, or with conditions resulting from cancer therapy, or treatment with immunosuppressive drugs, or as a consequence of AIDS. Most AIDS patients eventually contract Pneumocystis carinii pneumonia (PCP) and have accounted for the majority of the recent cases of this disease. Left untreated, PCP is almost always fatal.

The current method of treatment for P. carinii pneumonia is trimethoprim/sulfamethoxazole or pentamidine. Treatment with trimethoprim/sulfamethoxazole (TMP/SMZ) is associated with a high level of toxic side effects including rash, elevated liver function, nausea and vomiting, anemia, creatine elevation, and in extreme cases Stevens-Johnson syndrome. Side effects from TMP/SMZ are much more prevalent in patients with AIDS. Treatment with pentamidine is also associated with a high level of toxic side effects including renal failure, hepatotoxicity, hypoglycemia, hematologic abnormalities and pain or abscess at the injection site. The mortality due to treatment can reach 20 to 30 percent. An improved method for the treatment of P. carinii pneumonia in immune-compromised patients is greatly needed.

20

DE-A-2803581 describes a family of derivatives of the cyclic peptide echinocandin B, which are stated to have antimycotic activity.

EP-A-0032009, meanwhile, describes a related class of semi-synthetic cyclic peptides, which are stated to have antifungal activity.

25

DESCRIPTION OF THE INVENTION

The present invention is directed to the use of an anti-infection or infection controlling amount of a cyclohexapeptide compound for the preparation of a medicament, useful for treatment or prevention of Pneumocystis carinii infections in mammals.

30

The compounds within the scope of the present invention include a fungal metabolite as hereinafter detailed, known echinocandin type of antibiotics which are lipophilic cyclohexapeptides having a fatty acid side chain, simple derivatives of the echinocandin type antibiotics such as side chain dihydro- and tetrahydro- reduction products, ring ether and ester derivatives and semi-synthetic products in which the cyclohexapeptide ring is retained but in which the fatty acid side chain is replaced. Most of the compounds described in the literature are taught to be antifungal agents.

35

The compounds suitable for the practice of the present invention which include not only natural antibiotic compounds but also simple derivatives of natural antibiotic compounds and semi-synthetic compounds as are hereinafter fully detailed may be represented by the formula (I).

40

45

50

55



R' is H or OH;

R''' is H, OH, -O-



R''' is H or OH:

R'''' is H, CH_3 or CH_2CONH_2 ; and

R is

A)

• •




(B)





(C) when R' , R'' , R''' and R'''' are OH, and R''''' and R'''''' are CH_3 then

"



(ii)

or
(iii)

wherein

A is divalent oxygen, sulfur, sulfinyl or sulfonyl;

A' is divalent oxygen, sulfur, sulfinyl, sulfonyl or -NH-;

X is hydrogen, chloro, bromo, iodo, nitro, C₁-C₃ alkyl, hydroxy, C₁-C₃ alkoxy, mercapto, C₁, C₃ alkylthio, carbamyl or C₁-C₃ alkylcarbamyl;


X' is chloro, bromo or iodo;

R₁ is hydrogen, C₁-C₁₈alkyl or C₂-C₁₈alkenyl;W is C₁-C₁₀alkylene or C₂-C₁₀alkenylene;

m, n and p are 0 or 1, but if m is 0, n must be 0;


provided that the sum of the carbon atoms in the R₁ and W groups must be greater than 4 but cannot exceed 21; that when X is mercapto A and A' cannot be sulfinyl or sulfonyl; and that when A and A' are sulfinyl or sulfonyl, they must be in equal oxidation states;

(D) when R' is OH, R'' R''' and R'''' are H, then




wherein Y is a divalent aminoacyl radical of the formula

a)


wherein Z is C₁-C₁₀alkylene or C₅-C₆cycloalkylene

b)



wherein R₃ is hydroxyethyl, hydroxyethyl, mercaptomethyl, mercaptoethyl, methylthioethyl, 2-thienyl, 3-indolomethyl, phenyl, benzyl, halophenyl, halobenzyl, C₁-C₃ alkylphenyl, C₁-C₃ alkylbenzyl, C₁-C₃ alkylthiophenyl, C₁-C₃ alkylthiobenzyl, carbamylphenyl, carbamylbenzyl, C₁-C₃ alkylcarbamylphenyl or C₁-C₃ alkylcarbamylbenzyl;

(c)




wherein X² is hydrogen or halo.


The term "alkyl" is meant a univalent saturated straight chain or branched chain hydrocarbon radical. The term "alkenyl" is meant an univalent, unsaturated, straight chain or branched chain hydrocarbon radical containing not more than three double bonds. The term "halo" is meant chloro, bromo, fluoro and iodo.

The compounds useful in the method of treatment for or prevention of *Pneumocystis carinii* infections in mammals according to the present invention are generally white crystalline or powdery substances. They are generally of good solubility in alcohols such as methanol and ethanol, or poor solubility in water and in hydrocarbons such as hexane and of intermediate solubility in many common solvents. The solubility properties as well as other properties of many of the naturally occurring cyclohexapeptides are summarized in the CRC Handbook of Antibiotic Compounds, Vol IV, Part I, pp 355-367, CRC Press, Inc. Boca Raton, Florida, 1980.

An especially preferred compound is a novel fungal metabolite in which R', R'', R''' and R'''' are OH, R'''' is -CH₂CONH₂, R''''' is CH₃ and R is



represented by formula IA



25 The compound may be named 1-[4,5-dihydroxy-N²-(10,12-dimethyl-1-oxotetradecyl)-L-ornithine-5-(threo-3-hydroxy-1-glutamine)echinocandin B but hereinafter, for convenience, will be referred to as Compound IA. Especially preferred is the isomer named 1-[(4R,5R)-4,5-dihydroxy-N²-(10,12-dimethyl-1-oxo-tetradecyl)-L-ornithine-5-(threo-3-hydroxy-1-glutamine)echinocandin B. The preparation and properties of Compound IA are hereinafter described and are also the subject of copending applications Serial No. 105,795, filed October 7, 1987, corresponding to EP 311193, and Serial No. 105,797, filed October 7, 1987, corresponding to EP 311194, and are more fully described therein.

30 Compound IA, the preferred compound, is a white solid which may be characterized by the following physical properties:

a decomposition temperature of 206-214°C;

an empirical formula of C₅₁H₈₂N₈O₁₇ determined by high resolution FAB (Fast Atom Bombardment mass spectrometric measurement, calculated for C₅₁H₈₂N₈O₁₇ + Li = 1085.5958, found = 1085.6146);

35 an amino acid composition as determined by gas chromatogram/mass spectra of the trimethylsilyl derivative of the total acid hydrolysates of one equivalent each of threonine, hydroxyproline, methylhydroxyproline, and hydroxyglutamic acid.

¹H NMR Spectra in CD₃OD at 400 MHz as seen in Figure 1; and

40 ¹³C NMR chemical shifts obtained in CD₃OD at 100 MHz as follows: 11.20, 11.64, 19.75, 20.25, 20.78, 27.00, 28.07, 30.33, 30.37, 30.61, 30.76, 31.19, 31.29, 32.94, 34.83, 36.69, 38.10, 38.54, 39.07, 39.53, 45.93, 51.39, 53.01, 55.59, 56.31, 57.11, 58.35, 62.43, 68.18, 70.08, 70.55, 70.61, 71.26, 73.94, 75.72, 75.84, 76.86, 116.06(x2), 129.43(x2), 132.86, 158.22, 168.80, 172.16, 172.35, 172.40, 173.12, 174.24, 175.47, 176.88 ppm.

45 The compound is soluble in a variety of organic solvent such as methanol, ethanol, dimethylformamide, dimethyl sulfoxide, and the like.

The production of Compound IA, a novel compound, is described hereinafter in detail. The other compounds useful in the present invention may be prepared in a similar manner by substituting an appropriate microorganism or by other procedures as subsequently described.

50 Compound IA may be produced by cultivating a strain of microorganism designated MF 5171 in the culture collection of Merck & Co., Rahway, N.J. and identified as *Zalerion arboricola* and recovering said agent from the culture medium. A sample of the culture capable of producing the Compound IA has been deposited under the Budapest Treaty in the Culture Collection of the American Type Culture Collection at 12301 Parklawn Drive, Rockville, MD 20852. The sample has been assigned the accession number ATCC 20868.

The colonial and morphological description of ATCC 20868 are set forth below:

55 A. Morphological description

Globose, approximately 6.0 microns in diameter, thick-walled, dark coloured structures similar to chlamy-

dospores develop along mycelium and often appear to be intercalary. These structures appear to divide, forming multi-celled groups of 4-8 cells which do not break up readily. On some media, strands of mycelia cluster together, forming rope-like structures and the multi-celled groups form large clusters as if held together by mucilaginous material.

B. Colonial description

1. Czapek-Dox agar

Colonies are slow growing, growth not extensive, flat with irregular edges. Mycelium is black, shiny; surface becomes dull, granular-appearing, black to greenish-brown as colony ages.

2. Corn agar

Colonies are slow-growing. Growth is not extensive. Mycelium is black, shiny surface becoming powdery, dull black as colony ages.

3. Potato-dextrose agar, Sabouraud maltose agar and yeast extract-malt extract-dextrose agar.

Colonies are slow growing. Growth is not extensive, is slightly raised in the center, radiately furrowed irregular edges except on Sabouraud-maltose agar where the edges are hyaline and regular. Mycelium is black, shiny; becomes dull powder, black as colony ages.

Although the invention is discussed hereinbelow principally with respect to the specific strain, it is well-known in the art that the properties of microorganisms may be varied naturally and artificially. Thus, all strains of the genus ATCC 20868 including varieties and mutants, whether obtained by natural selection, produced by the action of mutating agents such as ionizing radiation or ultraviolet irradiation, or by the action of chemical mutagens such as nitrosoguanidine, are contemplated to be within the scope of this invention.

Compound IA may be produced in a form adaptable for drug use by cultivating the strain ATCC 20868 of Zalerion arboricola in a nutrient medium until a substantial amount of antibiotic activity is detected in the culture medium, generally in the mycelium portion thereof, and thereafter recovering the active component from the fermentation mycelium in a suitable solvent, concentrating the solution of active component, and then subjected the concentrated material to chromatographic separation.

The fermentation is carried out in a medium containing sources of carbon and nitrogen assimilable by the microorganisms and generally low levels of inorganic salts. In addition, the medium may be supplemented with trace metals, although if complex sources of carbon and nitrogen are employed, they are usually present in the complex sources.

The sources of carbon include glycerol, sugars, starches and other carbohydrates or carbohydrate derivatives such as dextran, cerelose, as well as complex nutrients such as oat flour, corn meal, millet and corn. The exact quantity of the carbon source which is utilized in the medium will depend, in part, upon the other ingredients in the medium, but it is usually found that an amount of carbohydrate between 0.5 and 5 percent by weight of the medium is satisfactory. These carbon sources can be used individually or several such carbon sources may be combined in the same medium.

The sources of nitrogen include amino acids such as glycine, arginine, threonine, methionine and the like as well as complex sources such as yeast hydrolysates, yeast autolysates, yeast cells, tomato paste, soybean meal, casein hydrolysates, yeast extracts, corn steep liquors, distillers solubles, cottonseed meal, and meat extract. The various sources of nitrogen can be used alone or in combination in amounts ranging from 0.2 to 90 percent by weight of the medium.

Among the nutrient inorganic salts, which can be incorporated in the culture media are the customary salts capable of yielding sodium, potassium, magnesium, ammonium, calcium, phosphate, sulfate, chloride, or carbonate. Also included are trace metals such as cobalt, manganese, iron, molybdenum, zinc, and cadmium.

The media suitable for carrying out the fermentation may be solid or liquid.

Solid media may have a millet, corn, oats, soy bean or wheat base. One medium having a millet base is Medium 2 in Example A. Other representative solid media include the following:

	<u>Media</u>	Weight or Volume Per <u>250 ml Flask</u>
5		
	<u>Medium A</u>	
	Corn (cracked)	10.0 g
10	Yeast extract	0.5 g
	Sodium tartrate	0.1 g
	Monosodium glutamate	0.1 g
	Corn oil	0.1 ml
15	Ferrous sulfate $\cdot 7\text{H}_2\text{O}$	0.01 g
	Water	15-20 ml
	<u>Medium B</u>	
20	Millet	15 g
	Yeast extract	0.5 g
	Sodium tartrate	0.1 g
	Ferric sulfate $\cdot 7\text{H}_2\text{O}$	0.01 g
25	Sucrose	0.5 g
	Alfalfa	0.5 g
	Corn oil	0.1 ml
30	Water	15 ml
	<u>Medium C</u>	
35	Millet	15 g
	Yeast extract	0.5 g
	Sodium tartrate	0.1 g
	Ferric sulfate $\cdot 7\text{H}_2\text{O}$	0.01 g
40	Silica gel	0.5 g
	Alfalfa	0.5 g
	Monosodium glutamate	0.1 g
45	Corn oil	0.1 ml
	Water	15 ml
50		
55		

Medium D

	Millet	15 g
5	Yeast extract	0.5 g
	Sodium tartrate	0.1 g
	Ferric sulfate•7H ₂ O	0.01 g

10

In addition, media may be prepared by substituting wheat, barley, oats or soy bean for the millet or corn above.

15

Liquid media also may be employed. Fermentation on a larger scale is generally more conveniently carried out employing a liquid medium. It has been found that although conventional liquid media may be employed to obtain Compound IA, such media have not been found to be suitable for obtaining good yields of the desired antibiotic. However, by incorporating from about 6 to 9 percent by weight of glycerol, it has been found that good yields of the desired antibiotic compound may be obtained. Suitable liquid media include:

20

Medium E

	Glycerol	85 g
	Pectin	10 g
25	Peanut Meal	4 g
	Peptonized Milk	4 g
	Tomato Paste	4 g
	Corn Steep	4 g
30	Lard Water	4 g
	Glycine	2 g
	KH ₂ PO ₄	2 g
35	Distilled Water to	1000 ml

pH 7.0

40

Medium F

	Dextrose	10 g
	Glycerol	10 ml
	Soy Flour	4 g
45	Peptonized milk	4 g
	Tomato paste	4 g
	Lard water	4 g
50	Potassium dihydrogen phosphate	2 g
	Cobalt chloride hexahydrate	0.01 g
	Distilled Water to	1000 ml

pH 7.0

55

For producing Compound IA, a fermentation medium containing ATCC 20868 is prepared by inoculating spores or mycelia of the antibiotic-producing organism into a suitable medium and then cultivating under aerobic conditions.

The procedure generally is first to inoculate a preserved source of culture from an agar slant containing nutrient medium into a nutrient seed-producing medium and to obtain, preferably through a two step procedure, growth of the organisms which serve as seeds in the production of the antipneumocystis agent.

In this process, a slant section of a preserved culture of ATCC 20868 is inoculated into an appropriate liquid nutrient seed medium of pH in the range 5 to 8.1, optimally 6 to 7.5, and the flasks incubated with or without agitation at temperatures in the range of from about 15°C to about 30°C, preferably 20° to 28°C. Agitation when employed, may be up to 400 rpm, preferably, about 200 to 220 rpm. The incubation is carried out over a period of from 2 to 30 days, preferably 2 to 14 days. When growth is abundant, usually between 2 and 5 days, the culture growth may be used to inoculate the production medium for the production of the antipneumocystis agent. Preferably however, a second stage fermentation is carried out, inoculating with a portion of the culture growth and then employing similar conditions but generally with shortened incubation period of about 1 to 3 days. The growth then is employed to inoculate the production medium.

The fermentation production medium inoculated with the culture growth is incubated for 3 to 30 days, usually 7 to 14 days with or without agitation. The fermentation may be conducted aerobically at temperatures ranging from about 20°C to about 40°C. For optimum results, it is most convenient to conduct these fermentations at a temperature in the range of from about 24°C to about 30°C. Temperatures of about 24°-28°C are most preferred. The pH of the nutrient medium suitable for producing the instant compounds can vary from about 5.0 to 8.5 with a preferred range of from about 6.0 to 7.5. After the appropriate period for the production of the desired compound or compounds, the latter is recovered from the fermentation medium as hereinafter more fully described.

The active material may be recovered from the fermentation medium by the steps comprising

- (1) adding alcohol to said medium, stirring and filtering to recover the active component in the resulting aqueous alcoholic solution;
- (2) concentrating the aqueous alcoholic solution to a small volume of primarily aqueous solution;
- (3) intimately contacting the resulting concentrated alcoholic aqueous solution with a water-immiscible organic solvent to extract or partition the active component thereinto and concentrating, or subjecting the concentrated solution to HP-20 adsorption and elution; then
- (4) subjecting the material recovered in Step (3) to at least one chromatographic separation, wherein in each chromatographic separation, the active component from the eluates exhibiting activity against Candida albicans are combined and concentrated to recover Compound IA.

The exact steps may vary somewhat depending on whether the fermentation had been carried out in liquid or solid medium, what solvent is employed and what adsorbent or combination of adsorbents is employed.

When the fermentation is carried out on solid medium, the first step may be carried out by adding an alcoholic solvent to the fermentation medium, thoroughly mixing, then filtering, recovering and concentrating the aqueous alcohol filtrate. Preferably, the concentrated filtrate is first back-extracted or washed with a lower aliphatic hydrocarbon solvent such as hexane or other alkane to remove alkane soluble impurities. The alkane washed filtrate may be extracted or partitioned with a water-immiscible oxygenated organic solvent and the resulting solution concentrated, then placed onto a column for at least one, generally several chromatographic separation steps. Suitable columns are silica gel, reverse phase and SEPHADEX LH-20.

When the fermentation is carried out in a liquid medium, in one method, the mycelial solids are filtered and recovered from the fermentation medium. Alcohol is added to the mycelial cake, and the mycelial solid thoroughly mixed with the alcohol, filtered, and the filtrate collected and concentrated. In an alternative method, the whole broth can be extracted by the addition of one volume of alkanol, preferably methanol, and filtered to remove solid impurities. The alkanol extract is then adsorbed on DIAION SP-207 or other commercially available styrene-divinylbenzene copolymer. A second dilution/HP-20 adsorption/elution step is utilized to concentrate the sample in preparation for chromatographic separations. Sometimes, a third dilution/HP-20 adsorption/elution step may be desirable for volume reduction.

The alcoholic solvent to be employed in the initial extraction of the active agent from the solid nutrient medium or from the mycelial pad may be any of the lower alcohols such as methanol, ethanol, isopropanol, and the like. Methanol is preferred.

The water-immiscible non-polar organic solvent useful for extracting or partitioning the active agent from the alkanol or methanol solution are esters, such as ethyl acetate, isopropyl acetate, butyl acetate, or ketones, such as methyl ethyl ketone. Lower aliphatic esters are preferred.

The chromatographic separation may be carried out by employing conventional column chromatography with non-ionic resin or by high performance liquid chromatography employing reverse phase resin. The fractions containing the antibiotic Compound IA may be detected by bioautography using Candida albicans. Generally, more than one chromatographic separation steps are employed. In a most preferred procedure, one or more separations are carried out employing column chromatography and a final separation is carried out em-

ploying high performance liquid chromatography (HPLC) with C₁₈ reverse phase resin.

When conventional column chromatography is employed for chromatographic separations, silica gel is the preferred adsorbent. Usually more than one chromatographic separation is required. Silica gel may be used in all the separations while employing different eluting agents. However, it may be combined advantageously with the use of a different adsorbent such as a dextran gel sold under the trade name of SEPHADEX LH-20 (Pharmacia). Other adsorbents such as alumina, styrene-divinylbenzene copolymers available commercially as DIAION HP-20, HP-30, HP-40, SP-207 (Mitsubishi Chemical Industries, Ltd.) and AMBERLITE XAD-2, XAD-4, XAD-16 (Rohm and Haas Co.) also may be employed.

In the fractionation and recovery of the active component by chromatography on silica gel, ester/alcohol mixtures with increasing concentration of alcohol provide good separations. A mixture of ethyl acetate and methanol has been found to be especially useful. These may be employed in isocratic, step gradient or continuous gradient systems. When a dextran adsorbent such as SEPHADEX LH-20, is employed, a chlorohydrocarbon/hydrocarbon/alcohol solvent system may be employed. A mixture of methylene chloride/hexane/methanol has been found to be especially useful.

In carrying out the HPLC separation, the alcohol solution containing material recovered from the conventional chromatography is concentrated and the residue dissolved in methanol and placed on a column packed with commercial reverse phase resin or on a column filled with silica gel/C₁₈ reverse phase resin prepared as amply reported in the literature. Alternatively, the alcohol solution may be diluted to 50 percent with water and pumped onto the column. The column is operated using methane/water (1:1 or optionally other ratios) at 800-2000 psi [(5.5-13.8)x10⁶ Pa] which produces a flow rate of about 20 ml/min. Separation is monitored at 210 nm.

The fractions are assayed for activity with Candida albicans. The product is recovered from the chromatographic procedures by combining the active fractions and concentrating under reduced pressure. The product then may be purified by conventional chromatography or preparative HPLC.

Other natural products which are embraced by the present invention include those which are described in the literature as echinocandins, aculeacins, mulundocandins, athlestin (also referred to as an echinocandin T), sporiofungin or by number designations. The structures in some cases are not completely identified but the compounds are known to be neutral cyclohexapeptides with fatty acid side chain.

The echinocandins include echinocandin-A (also known as A-32204-A) produced by Aspergillus nidulans-echinulatus (Swiss patent 568,386); echinocandin-B (also known as A-32204-B and SF-7810-F) produced by Asp. nidulans (Swiss 568,386) and Asp. rugulosus (Helv. Chim. Acta 62 11252 (1979); German 2,549,127; Belg. 834,289); echinocandin-C (also known as SL-7810-FII) produced by Asp. rugulosus (Ger. 2,549,127; Belg. 834,289); echinocandin-D (also known as SL-7810-FIII) produced by Asp. rugulosus (Ger. 2,549,127; Belg. 834,289).

Echinocandins B, C and D may be produced together on the fermentations of a strain of Aspergillus rugulosus. The major metabolite in the fermentation of Aspergillus rugulosus is echinocandin B, having a molecular formula of C₅₂H₈₁N₇O₁₆. Echinocandin C and D which are the minor metabolites differ in certain of the amino acids making up the cyclic peptide ring. Thus, echinocandin C, having a molecular formula of C₅₂H₈₁N₇O₁₅, contains 3-hydroxyhomotyrosine in place of the 3,4-dihydroxyhomotyrosine present in echinocandin B. Echinocandin D, having a molecular formula of C₅₂H₈₁N₇O₁₃, contains 3-hydroxyhomotyrosine and ornithine instead of 3,4-dihydroxyhomotyrosine and 4,5-dihydroxyornithine.

Echinocandin B also may be produced by the cultivation of Emericella nidulans as described in the working example.

The aculeacins include aculeacin-A, aculeacin-B, aculeacin-C, aculeacin-D, aculeacin-E, aculeacin-F, aculeacin-G, aculeacin-AA", aculeacin-DA", aculeacin-A G", aculeacin-Aα and Ay, and aculeacin-Dα and Dy. The aculeacins may be produced by the fermentation of Aspergillus aculeatus (either liquid or solid culture) (preferably M4214 FERM-P 2324) thereafter extracting the mycelium with a water miscible organic solvent concentrating and purifying by chromatography as described in U.S. Patent 3,978,210 and 4,212,858 German Patent 2,509,820 and Belgian Patent 826,393.

Another cyclohexapeptide antibiotic found in the literature is mulundocandin (R', R'', R''', R'''' are OH, R'''' is H, R is -CO(CH₂)₁₀-CH₃CH-CH₂CH₃) produced by Asp. sydowii No. Y-30462 (J. Antibiotics 40, 275 and 40, 281 (1987)). The antibiotic, which is present both in the culture filtrate and mycelium may be isolated from the ethyl acetate extract of a culture filtrate by silica gel column chromatography and droplet counter current chromatography as more fully described by Roy et al, J. Antibiotics 40, 275-280 (1987).

Still another echinocandin type cyclopeptide reported in the literature is sporiofungin (R', R'''' and R''' are OH, R'' is H, R''' is CH₂CONH₂, R'''' is H and R is a branched chain C₁₆ fatty acid radical,) which may be produced by cultivation of Cryptosporiopsis (WIPO publication, WO 82/00587) strain ATCC 20594 or NRRL 12192, on various conventional nutrient media as aerobic surface culture or immersion culture, and the mycelium sepa-


rated from the culture broth and recovered in a conventional manner such as by homogenizing the mycelium with methanol, extracting from the methanol with a water-immersible organic solvent such as ethyl acetate, removing the solvent and purifying by chromatography as more fully described in WO 82/00587.

Other echinocandin type cyclohexapeptides include X-73 produced by *Aspergillus rugulosus*, (Ind. J.B. Biochem. Z, 81 (1970)), athlestatin produced by *Aspergillus niger*, (Jap. J. Ant., 17 268 (1964), JP 66/12688; CA 65 19274; 70, 27607), S-31794-F-1 produced by *Acrophialophora limonisporea* (U.S. 4,173, 629; Ger. 2,628,965), A-30912-A or A-22082 produced by *Asp. nidulans* and *Asp. rugulosus* (U.S. 4,074,246; 4,074,245; 4,288,549; Ger. 2,643,485; CA 87, 20612) A-30912-B, A-30912-C, and A-30912-D, produced by *Asp. rugulosus* (U.S. 4,024,245).


Compounds which are modifications of natural products include those in which the side chain, i.e., R, when an unsaturated fatty acid, had been modified by reduction. This may be carried out catalytically, by using Pd on carbon at atmospheric pressure.

Compounds in which R''' is -O-Alkyl (C₁-C₆) or -O-benzyl may be prepared by stirring together the corresponding compound in which R''' is OH with the appropriate alkanol under acidic conditions at ambient temperature for several hours or overnight to obtain the ether in the mixture and thereafter preferably quenching the reaction with dilute bicarbonate and recovering it by conventional procedures. In addition to the preparation in the working example, the preparation for a number of these compounds is more fully described in GB 2,050,385A.

Compounds in which R''' is



wherein R¹ is hydrogen, alkyl or substituted alkyl as previously defined, R² is hydrogen or C₁-C₄ alkyl, or R¹ and R² together form -(CH₂)₂-Q-(CH₂)₂- wherein Q is a connecting group such as -CH₂-, -NH- or -N(lower alkyl)- may be prepared by reacting a compound in which R''' is OH with a compound of the formula



in the presence of an organic acid such as p-toluenesulfonic acid, methanesulfonic acid or mineral acid in an aprotic solvent such as dimethylformamide as more fully described in Belgian 851,310.

Semi-synthetic compounds in which the acyl side chain, i.e., R in above formula, has been modified from naturally occurring fatty acids to structurally distinctive acyl groups may be produced by first removing the fatty acid side chain and thereafter introducing a distinctive acyl group. The fatty acid side chain preferably is removed enzymatically. The enzyme which is useful for deacylation of the cyclohexapeptides to obtain a cyclohexapeptide nucleus is that produced by certain microorganisms of the family *Actinoplanaceae*, especially the microorganism *Actinoplanes utahensis* NRRL 12052 which is available from the Northern Regional Research Center, USDA, Agricultural Research Service, Peoria 111 61604, or as *A. utahensis* ATCC 14539 obtainable from the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852.

The enzyme may be produced by growing *Actinoplanaceae* at temperatures between about 25°C and 30°C and a pH of between about 5.0 and 8.0, preferably between 6.0 and 7.0, with agitation and aeration for from about 40 to about 60 hours in a culture medium containing (a) an assimilable carbon source such as sucrose, glucose or glycerol, (b) a nitrogen source such as peptone, urea or ammonium sulfate (c) a phosphate source such as a soluble phosphate salt and (d) growth promoting inorganic salts.


In the deacylation, the cyclohexapeptide compound or substrate containing the cyclohexapeptide is added to the culture of *Actinoplanaceae* after the culture has been incubated for at least 48 hours. After addition of the substrate, the incubation of the culture is continued for about 24 hours or longer over temperatures in the range of from about 25°C to about 30°C.

The course of the reaction may be monitored by *Candida albicans* assay. The starting cyclohexapeptide compound is active against *C. albicans* but the deacylated nucleus compound is biologically inactive.

The deacylated nucleus compound then may be employed in the preparation of semi-synthetic compounds. Conventional acylation methods may be employed. In one preferred method, a 2,4,5-trichlorophenyl

ester of the desired acid is reacted with the deacylated nucleus compound in an inert solvent such as dimethylformamide at room temperature for about 15 to 18 hours. This may be illustrated with a specific example as follows:

A deacylated nucleus of a compound the following structure wherein in formula (IB), E is H (Compound IBi)



may be prepared by first preparing a deacylating enzyme by the fermentation of *Actinoplanes utahensis* NRRL 12052 by first inoculating a slant with the organism, incubating the slant at 30°C for about 8 to 10 days and then using the slant growth to inoculate 50 milliliters of vegetative medium which is incubated at 30°C for about 72 hours with shaking. The incubated vegetative medium may be used to inoculate 400 milliliters of a second-stage vegetative medium having the same composition as the first stage vegetative medium and incubated at 30°C for about 48 hours with shaking. Thereafter, 800 milliliters of said medium may be used to inoculate 100 litres of production medium which is permitted to ferment at about 30°C for 42 hours with agitation and aeration to maintain dissolved oxygen level above 30 percent of air saturation at atmospheric pressure and to obtain therein the desired deacylating enzyme.

The medium for preparing the slant may be of the following composition: Baby oatmeal 60.0 g; yeast 2.5 g; K_2HPO_4 1.0 g; Czepek's mineral stock 5.0 ml; agar 25.0 g; deionized water q.s. to 1 liter. Czepek's mineral stock is of the following composition: $FeSO_4 \cdot 7H_2O$ (in 2 ml. conc. HCl) 2 g; KCl 100 g; $MgSO_4 \cdot 7H_2O$ 100 g; deionized water to 1 liter with final adjusted pH 6.7.

The medium suitable for both first stage and second stage vegetative medium may be of the following composition: Baby oatmeal 20.0 g; sucrose 20.0 g; yeast 2.5 g; dried grain 5.0 g; K_2HPO_4 1.0 g; Czepek's mineral stock 5.0 ml and deionized water to 1 liter, final pH 6.8.

The production medium may be of the following composition: Peanut meal 10.0 g; soluble meat peptone 5.0 g; sucrose 20.0 g; KH_2PO_4 0.5 g; K_2HPO_4 1.2 g; $MgSO_4 \cdot 7H_2O$ 0.25 g; tap water q.s. to 1 liter.

An alcohol solution of a cyclohexapeptide (IB) where E is stearoyl (tetrahydroechinocandin B) or palmitoyl (aculeacin A) is added to the production fermentation medium above-described and the deacylation monitored by assay against *Candida albicans*.

On completion of the deacylation, the fermentation broth containing the nucleus compound is filtered, passed through HP-20 resin to place the nucleus compound on the column and the compound recovered by elution with water/methanol (7/3) at a rate of 200-300 ml/minute. The eluate is monitored for the nucleus compound by treating with acid chloride and assaying for activity against *Candida albicans*. The eluate fractions showing activity are concentrated to obtain the nucleus compound (IBi).


The deacylated cyclohexapeptide compound thus obtained is purified by dissolving in water/acetonitrile/acetic acid/pyridine (96/2/1/1) and purified by reversed phase liquid chromatography (LICHROPREP RP-18) at a flow rate of about 60 milliliters per minute and monitoring the separation at 280 nm using a UV monitor. On the basis of the absorption at 280 nm, the fractions containing the desired compound are combined, concentrated under reduced pressure and employed for acylation or lyophilized for subsequent use.

The preparation of the deacylated cyclohexapeptide compound having the same nucleus from echinocan-

din type natural products is more fully described in U.S. 4,293,482. Similar preparations of other deacylated cyclohexapeptide compounds with similar nuclei may be found described in U.S. patents 4,173,629; 4,293,490; 4,299,763; 4,299,762; and 4,304,716.

The deacylated compounds may then be employed to produce novel and/or unusual acylated compounds.

The acylation of the compound of the deacylated nucleus, to produce a unique acyl derivative may be illustrated with a preparation of a compound in which E in formula (IB) is




(Compound IBii). Such a compound would also be illustrated of a compound of formula (I) in which R is a radical defined under (C)(i).

In the acylation, the acyl group is preferably introduced by means of a trichlorophenyl ester. Thus, first, the trichlorophenyl ester may be prepared by treating the side chain acid with 2,4,5-trichlorophenol in the presence of a coupling agent such as N,N'-dicyclohexylcarbodiimide in an inert solvent such as methylene chloride. About 1.1 mole of the 2,4,5-trichlorophenol and 1.0 mole of the coupling agent is used for each mole of the alkoxybenzoic acid. The mixture is stirred at room temperature for 15 to 18 hours to obtain the ester which may be recovered by filtering the solid and evaporating the filtrate under reduced pressure, then recrystallizing the residue.

The ester thus prepared is added to a solution of the nucleus compound in dimethylformamide and stirred for about 18 hours and then the solvent evaporated off. The residue is washed and then chromatographed on silica gel using ethyl acetate-methanol (3/2) as eluant to obtain the desired octyloxybenzoyl derivative which may be represented in formula IB with the octyloxybenzoyl group as E. Such a compound may be named 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithine]echinocandin B (Compound IBii).

In addition to Compound IA, other preferred compounds include Compound IBii; echinocandin B (R', R'', R''', R'''' are OH; R''''' and R'''''' are CH₃; R is



tetrahydroechinocandin B (R', R'', R''', R'''' are OH; R''''' and R'''''' are CH₃; R is

0

In a representative study employing Compound IA in rats, nine male Sprague-Dawley rats weighing approximately 200 grams each, were immunosuppressed by the addition of dexamethasone to the drinking water (2.0 mg/liter) for six weeks to induce the development of *P. carinii* infections. To enhance the infection the rats were also maintained on a low protein diet. At the beginning of the seventh week the rats were divided into three groups of three rats each. All three groups continued to receive dexamethasone in the drinking water and low protein diet throughout the remainder of the study. The rats in Group I were kept as untreated infected controls, those in Group II were injected intraperitoneally twice daily for two weeks with 0.5 ml of sterile dH₂O containing 2 mg of Compound IA, and those in Group III were treated with trimethoprim-sulfamethoxazole (TMP-SMZ) in the drinking water (208 mg TMP and 1.040 g of SMZ/liter) for two weeks, a known treatment for *P. carinii* infections. Two rats died early in the experiment. One was from group II (rat 72A) and the other from group III (rat 75A). It was not determined whether they died from *Pneumocystis pneumonia*.

At the end of the two week treatment period (a total of eight weeks of immunosuppression) the animals were sacrificed and the lung tissue removed. The lungs from each animal were weighed, and then processed to determine the number of cysts and parasite nuclei for each animal. The results of this experiment are shown in Table I.

TABLE I
IN VIVO TESTING OF COMPOUND IA & TMP-SMZ
FOR ACTIVITY AGAINST *P. CARINII*

RAT	TOTAL CYSTS			TOTAL NUCLEI			LUNG WT. (gms)	CYST SCORE
	#	per lung	per gm lung	per lung	per gm lung			
GROUP I (control)	714	3.2x10 ⁸	1.0x10 ⁸	5.8x10 ⁹	1.8x10 ⁹	3.16	4	4
	74A	1.5x10 ⁹	4.5x10 ⁸	1.4x10 ¹⁰	4.2x10 ⁹	3.32	4+	4+
	74B	7.5x10 ⁸	2.1x10 ⁸	1.4x10 ¹⁰	3.9x10 ⁹	3.56	4+	4+
GROUP II (COMPOUND IA)	71B	N.D.*	N.D.*	2.1x10 ⁹	7.4x10 ⁸	2.83	0	0
	72B	N.D.*	N.D.*	1.0x10 ⁹	7.1x10 ⁸	1.40	0	0
GROUP III (TMP-SMZ)	73B	6.0x10 ⁶	5.4x10 ⁶	2.0x10 ⁸	1.8x10 ⁸	1.11	1	1
	75B	5.4x10 ⁶	2.3x10 ⁶	1.6x10 ⁹	7.0x10 ⁸	2.28	1	1

*N.D. = No cysts were detected in 50 fields, indicating that there were less than 2×10^5 (limits of resolution).

These results demonstrate the Compound IA has efficacy against *P. carinii* infections. No cysts could be found in the lungs of either rat treated with Compound IA as compared to levels ranging from 3.2×10^8 - 1.5×10^9 in the control animals. There was also a reduction in the number of parasite nuclei (1.0 - 2.1×10^9) in these animals as compared to controls (5.8×10^9 - 1.4×10^{10}). These results are superior to those seen in animals treated with TMP-SMZ, a known treatment for *P. carinii* infections, since the TMP-SMZ rats had detectable levels of *P. carinii* cysts.

A similar rat study was conducted to determine the antipneumocystis activity of 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ii ornithine]echinocandin B (Compound IBii). After 6 weeks of immunosuppression the rats were injected twice daily for 7 days with vehicle alone. Compound IBii (2 mg/kg) or Compound IA (2 mg/kg) during which time immunosuppression was continued. At the end of the treatment period the animals were sacrificed and the number of cysts per lung determined. Compound IA reduced the number of cysts by >99% while Compound IBii reduced the cyst counts by approximately 84% (compared to the vehicle control, 10% DMSO).

Studies were also carried out with mice and still other compounds within Formula I. Representatives of the method used in the studies carried out with mice and with other compounds is as follows:

Male C3H/HeN mice, weighing 22-24 grams each, were immunosuppressed by the addition of dexamethasone to the drinking water for six weeks to induce the development of *P. carinii* infections. To enhance the infections the mice were also maintained on a low protein diet. At the beginning of the seventh week, the mice were randomly divided into groups of 6. All groups continued to receive dexamethasone in the drinking water and low protein diet for the remainder of the study. Mice in each group were injected intraperitoneally twice daily with 0.5 ml of a 10% DMSO solution as a vehicle control or the same cocktail containing drug to achieve a dose of 2 mg/kg. This procedure was continued for one week.

At the end of the treatment period (a total of seven weeks of immunosuppression) the animals were sacrificed and the lung tissue removed. The tissue was then processed to determine the number of cysts for each animal.

The results for the various compounds are seen in Table II.

TABLE II

<u>Compound</u>	<u>Dosage</u>	<u>% Reduction of Cysts</u>
Tetrahydroechinocandin B (TEB)	2mg/kg.	99%
O-Methyl-TEB	2mg/kg	>99%
O-Benzyl-TEB	2mg/kg	99%
Echinocandin B	2mg/kg	>99%
Compound IA	2mg/kg	>99%

From the foregoing test results and from known dosage ranges for TMP-SMZ as applied to man, it is determined that generally from about 0.5 to about 20.0 mg/kg of body weight of the cyclohexapeptide anti-pneumocystis agent (Compound I) may be employed while considering the patients' health, weight, age and other factors which influence response to a drug as well as whether it is to be applied to a human patient or to an animal. These amounts, when expressed as doses suitable for human beings, are in the range of from about 35 mg to about 1500 mg daily preferably by parenteral administration.

The outstanding properties are most effectively utilized when the compound is formulated into novel pharmaceutical compositions with a pharmaceutically acceptable carrier according to conventional pharmaceutical compounding techniques.

The novel compositions contain at least 1% by weight of the active compound. In preparing the compositions, Compound I is intimately admixed with any of the usual pharmaceutically acceptable carrier.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), pulmonary (nasal or buccal inhalation), nasal administration, or insufflation.

The cyclohexapeptide anti-pneumocystis agent, Compound I, is preferably formulated into compositions for injection and may be presented in unit dosage form in ampoules or in multidose containers, if necessary with an added preservative. The compositions may also take such forms as suspensions, solutions or emulsions in oil or in vehicles such as 20/80 ethanol/propylene glycol or 20-35% polyethylene glycol 300 and may contain formulating agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active

ingredients may be in powder form for reconstituting with a suitable vehicle prior to administration or for use in insufflation.

The term "unit dosage form" as used in the specification and claims refer to physically discrete units, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the pharmaceutical carrier. Examples of such unit dosage forms measured units in ampoules or in multidose containers and the like. A unit dosage of the present invention will generally contain from 100 to 1000 milligrams of the cyclohexapeptide anti-pneumocystis agent.

Any suitable route of administration may be employed for providing a patient with an effective dosage of a compound of Formula I. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, disperions, suspensions, solutions, capsules, creams, ointments, aerosols, powders for insufflations and the like.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), pulmonary (nasal or buccal inhalation), nasal administration, or insufflation.

For administration by inhalation, the compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The compounds may also be delivered as powders which may be formulated and the powder compositions may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery system for inhalation is a metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of Compound I in suitable propellants, such as fluorocarbons or hydrocarbons.

Because of the difficult solubility of the compound in pharmaceutically acceptable liquid carriers and because of desirability to directly treat lung and bronchi, aerosol administration is a preferred method of administration. Insufflation is also a desirable method, especially where infection may have spread to ears and other body cavities.

Alternatively, parenteral administration may be employed using drip intravenous administration. In such method, the drug may be solubilized in alcohol/propylene glycol or polyethylene glycol compositions. Intraperitoneal injection also may be employed.

The following examples illustrate the preparation of Compound IA and other lipophilic cyclohexapeptide compound of Formula I and the compositions useful in the present invention of controlling Pneumocystis carinii.

PREPARATION OF COMPOUND IA

EXAMPLE A

Fermentation

A frozen culture in glycerol of Isolate 2 of Culture 8525-307P originally isolated from water, identifiable as Zalerion Arboricola, ATCC 20868, and maintained in the Merck culture collection was employed in the fermentation.

A 2 milliliter portion of the frozen culture was defrosted and aseptically transferred to a 250 milliliter unbaffled Erlenmeyer flask containing 54 milliliters of Medium 1. Medium 1, after inoculation, was incubated at 28°C with rotary agitation (220 rpm, 2" throw shaker) for three days. At the end of this period, 2.0 milliliters of the growth medium were aseptically transferred to each of several unbaffled 250 milliliter Erlenmeyer flasks containing Medium 1. The inoculated flasks were incubated at 28°C for 2 days.

12.5 milliliters of the mature seed broth were inoculated into five production flasks containing Medium 2, and incubated at 25°C for seven days under static conditions to obtain an antibiotic compound on the fermentation medium.

The media employed in the foregoing fermentation were:

MEDIUM 1 (KF Seed Medium)

5	Corn Steep Liquor	5 g
	Tomato Paste	40 g
	Oat Flour	10 g
	Glucose	10 g
10	Trace Elements Mix	10 ml
	Distilled Water	to 1000 ml
	pH 6.8	

15

Trace Elements Mix:

20	FeSO ₄ ·7H ₂ O	1 g
	MnSO ₄ ·4H ₂ O	1 g
	CuCl ₂ ·2H ₂ O	25 mg
	CaCl ₂	100 mg
25	H ₃ BO ₃	56 mg
	(NH ₄) ₆ MoO ₂ ·4H ₂ O	19 mg
	ZnSO ₄ ·7H ₂ O	200 mg
30	Distilled Water	to 1000 ml

MEDIUM 2 (F204 Solid Medium)

35

	Amount/flask
Millet Base	15 g
40 Yeast Extract	.5 g
Sodium tartrate	.1 g
Ferrous Sulfate Crystals	.01 g
45 Monosodium Glutamic Acid	.1 g
Corn Oil	.1 ml

50 Isolation

Five hundred milliliters of methanol were added to each of the five 2-liter flasks of solid phase fermentation. The contents of the flask were then combined and stirred to extract methanol-soluble material and the mixture then filtered. The spent cake was stirred with an additional 2500 milliliters of methanol to further extract methanol soluble material and the mixture then filtered.

55

The filtrate and wash were combined and concentrated to 500 milliliters.

The aqueous methanolic concentrate thus obtained then was extracted with two 500 milliliter portions of ethyl acetate. The spent aqueous solution was placed onto a DIAION HP-20 column to adsorb the active ma-

terial thereon and the latter then eluted therefrom with methanol. The eluates were combined with the previously obtained ethyl acetate extracts and the combined ethyl acetate solutions were concentrated to dryness. The residue was chromatographed on 200 milliliters of SEPHADEX LH-20 using 5:5:2 methylene chloride/hexane/methanol as eluant.

5 The fractions active as determined by Candida albicans were combined and chromatographed on 200 milliliters of silica gel (EM Science, KEISELGEL 60, 230-400 mesh) using a step gradient elution with ethyl acetate/methanol. The active fractions from this chromatography were combined, concentrated and chromatographed on silica gel using a 75:25 ethyl acetate/methanol isocratic system.

10 The active portions from this chromatography were then combined and placed on 100 milliliters of SEPHADEX LH-20 using methanol as eluting solvent. The eluate, after vaporization of the solvent, yielded 95 milligrams of a purified compound. The compound was a white solid having a ¹H NMR spectrum as seen in Figure 1.

EXAMPLE B

15

Fermentation

In a manner similar to that described in Example A, the contents of one frozen vial of ATCC 20868 from the Merck culture collection were defrosted and aseptically transferred to a 250 milliliter un baffled flask containing 54 milliliters of KF medium (Medium 1) containing 0.4 percent agar. The modified Medium 1, after inoculation, was incubated at 28°C with 220 rpm agitation for 48 hours. At the end of this period, 10 milliliters of the growth medium was transferred to a 2-liter un baffled flask containing 500 milliliters of KF medium containing 0.4 percent agar. After inoculation, the resulting medium was incubated for 24 hours at 28°C with 220 rpm agitation.

25 Twenty 2-liter flasks each containing 120 grams of Medium 2 and 120 milliliters of a stock solution consisting of

Yeast extract	5 parts by weight
Sodium tartrate	1 part by weight
Ferrous sulfate crystals	0.1 parts by weight
30 Monosodium glutamic acid	1 part by weight
Corn oil	1 part by weight

were autoclaved for 20 minutes at 122°C and then reautoclaved with 80 milliliters of water for another 20 minutes at 122°C. The flasks were allowed to cool, then inoculated with 20 milliliters of seed medium prepared as above described, and the inoculated flasks incubated at 25°C under static conditions for 14 days.

35

Isolation

To each of nineteen 2-liter solid fermentation flasks was added 1 liter of methanol and the contents combined, stirred and filtered. The spent cake was extracted twice with 6 liters of methanol. The aqueous methanol 40 filtrates were then concentrated and the concentrate extracted twice with 3 liters of ethyl acetate. The ethyl acetate extracts were combined, dried and concentrated to about 100 milliliters.

The concentrate was then coated on silica gel by adding 100 milliliters of methanol and 100 milliliters of silica gel thereto, intimately contacting the components and then removing the solvent on a rotary evaporator. The dried silica gel was then applied to a column of 500 milliliters of silica gel, the column washed with ethyl 45 acetate to remove impurities and eluted with 9:1 ethyl acetate/methanol. The eluates containing antibiotic material testing positive against Candida albicans were recovered and combined.

The antibiotic rich cut from the silica gel chromatography was dissolved in 200 milliliters of 10:10:1 methylene chloride/hexane/methanol and the resulting solution combined with 40 milliliters of SEPHADEX LH-20 (which previously had been prepared by soaking overnight in methanol followed by washing twice with 200 50 milliliters of methylene chloride/hexane/methanol). After a few minutes, the supernatant was removed by filtration and the SEPHADEX LH-20 was washed with 200 milliliter of methylene chloride/hexane/methanol and then filtered. The filtrates were found not to contain the active constituent, and were discarded. The active constituent which had partitioned into the dextran SEPHADEX LH-20 beads was extracted therefrom by washing twice with 200 milliliters of methanol, and these methanol washes were combined and concentrated.

55 The methanol concentrate was next applied to 200 milliliters of silica gel and eluted with 75:25 ethyl acetate/methanol and the eluates combined and the solvent vaporized to obtain Compound IA. Compound IA is a white powder having a decomposition point of 206-214°C.

PREPARATION OF ECHINOCANDIN B DERIVATIVES

EXAMPLE C

5 Preparation of Echinocandin BA. Fermentation

10 Culture MF 5100 *Emericella nidulans* ATCC 20956 from the Merck culture collection was used for seed train development as follows: One frozen vial (about 2.5 ml) of MF 5100 was used to inoculate 54 mL of KF medium and cultured in 250 ml unbaffled Erlenmeyer flasks at 28°C and 220 rpm for 72 hours to obtain "B" stage culture. Five mL of "B" stage culture was used to inoculate 500 mL of KF medium in 2 liter unbaffled Erlenmeyer flask and incubated at 28°C and 220 rpm for 48 hours to obtain "C" stage cultures. Four "C" stage flasks were used to inoculate 160 liters of KF medium with trace element mix #2 in a "D" stage (300 liter stirred fermenter).

The three stages of seed train were used for "E" scale fermentation in an 800 liter fermenter. The fermentation was carried out for 23 hours at 28°C, back pressure 0.7 kg/cm², aeration with 0.300m (53 liters per minute) and agitation of 100 rpm.

20 The production "E" stage was inoculated with 5% seed from "D" stage in 5700 liters of a production medium of the following composition:

Glycerol	85 g/L
Pectin	10 g/L
Peanut Meal	4 g/L
Peptonized Meal	4 g/L
25 Peptonized Milk	4 g/L
Tomato Paste	4 g/L
Corn Steep Liquor	4 g/L
Lard Water	4 g/L
Glycine	2 g/L
30 Potassium phosphate monobasic	2 g/L
Polyglycol P-2000	1 mL/L

The medium was sterilized for 25 minutes at 123°C and the fermentation then carried out at temperature of 28°C, positive pressure of about 0.8 kg/cm², aeration 6000 liters per minute, agitation speed of 120 rpm; pH in the range 5.5-6.5 with dissolved oxygen maintained above 30 percent for about 36 hours.

35

B. Isolation

The whole broth from the above fermentation containing 355 grams of echinocandin B by HPLC assay (against reference standard echinocandin B) was extracted overnight with an equal volume of methanol. The diluted broth was centrifuged to remove broth solids and the centrifugate adsorbed on a 450 liter SP 207 column which had been equilibrated with 50:50 methanol/water. After a 1000 liter water wash and a 1000 liter 50:50 methanol/water wash, the column was eluted with methanol and the eluate concentrated to 189 liters and partitioned with methylene chloride. The aqueous methanol layer was adjusted to 50:50 methanol/water by the addition of water and adsorbed on a 100 liter HP-20 column prewashed with methanol and pre-equilibrated with 50:50 methanol/water. After a 50:50 methanol/water wash, the column was eluted with 80:20 methanol/water. The 80:20 eluate was adjusted to 50:50 with water, and then was adsorbed on a 27 liter SP207 column, preequilibrated as before. The column was then eluted with methanol.

45 A 19 g aliquot of SP 207 eluate was diluted to 50:50 methanol/water with water and adsorbed on a 3.9 liter Amicon preparative C-18 column. A gradient of acetonitrile/water (25:75) to 100 percent acetonitrile over a 75 minute period was used to elute the compound. Selected fraction after concentration and lyophilization from acetonitrile/water provided 16 grams of Echinocandin B of 87-91 percent purity.

50 The identity and purity of the foregoing echinocandin B was determined by HPLC and spectral comparison with a sample of Echinocandin B, of the properties below, previously prepared and purified, and its identity established by comparison with a published structure of Echinocandin B.

55

EXAMPLE D

Preparation of Tetrahydroechinocandin B

5 5.0 Grams of Echinocandin B (EB) of about 87 percent purity (HPLC) was dissolved in 150 milliliters of absolute ethanol and was added to 10% Pd-C in 80 milliliters of ethanol which had been prereduced under 1 atmosphere of hydrogen for 20 minutes. After the addition, the mixture was stirred rapidly under 1 atmosphere of hydrogen for 4 hours. Analysis by HPLC using CH₃CN/H₂O (81/19), on a C8 ZORBAX column at flow rate 1 ml/min at 30°C showed that the reaction was virtually complete at this time.

10 The reduction mixture was filtered through a Celite pad, the pad washed with ethanol and the mixture was concentrated in vacuo to obtain a solid. The solid was purified in ten 0.5 gram batches dissolved in 5 milliliters of CH₃CN/H₂O (81/19). The HPLC was carried out on C8 ZORBAX 1 inch column and eluted with 80% methanol/20% water at a flow rate of 15 ml/min. A total yield of 3.15 grams (79%) of tetrahydroechinocandin B was obtained.

15

EXAMPLE E

Preparation of O-Methyltetrahydroechinocandin B

20 55 Milligrams of tetrahydroechinocandin B was dissolved in 4 milliliters of methanol. To the solution was added 4.0 milligrams of p-toluenesulfonic acid and the mixture stirred at room temperature for 3 hours. HPLC analysis indicated that the reaction was complete at this time.

The reaction mixture was quenched with 1N sodium bicarbonate solution and washed twice with saturated sodium chloride solution and then diluted with 200 milliliters of isopropanol/chloroform (2/7).

25 The organic layers were dried over sodium sulfate, filtered and the solvent vaporized to obtain a near white solid as residue. The latter was purified first with methanol, and then with H₂O/CH₃CN (2:3) and chromatography by MPLC in H₂O/CH₃CN (2:3) on a C8 LOBAR size A column in 4 milliliter fractions. The compound was obtained in a yield of 42.4 mg. (76%)

30 EXAMPLE F

Preparation of O-Benzyltetrahydroechinocandin B

35 50 Milligrams of tetrahydroechinocandin B was suspended in 1 milliliter of tetrahydrofuran. To it was added 97 milliliters of anhydrous benzyl alcohol and the resulting mixture stirred until all was in complete solution. To the resulting solution was added 1 mg. of p-toluenesulfonic acid monohydrate and the mixture then stirred at room temperature for 3 hours whereupon a HPLC analysis showed that the reaction was complete. The reaction mixture was diluted with 1N sodium bicarbonate solution and then with isopropyl alcohol/chloroform (3:7). The organic solution was washed twice with water, dried and evaporated to recover the product as residue.

40

The following examples illustrate novel compositions useful in the practice of the present invention.

Aerosol compositions may be prepared having the following formulations:

EXAMPLE I

45

	<u>Per Canister</u>
Compound Ia	24 mg
50 Lecithin, NF Liquid Concentrate	1.2 mg
Trichlorofluoromethane	4.025 g
Dichlorodifluoromethane	12.15 g

55

EXAMPLE II

	<u>Per Canister</u>
5 Echinocandin B	24 mg
Oleic acid	1.2 mg
Trichlorofluoromethane	4.025 g
10 Dichlorodifluoromethane	12.15 g

EXAMPLE III

	<u>Per Canister</u>
15 O-Methyltetrahydroechinocandin B	24 mg
Sorbitan trioleate	1.2 mg
20 Trichlorofluoromethane	4.025 g
Dichlorodifluoromethane	12.15 g

25 An injectible suspension (1M) may be prepared as follows:

EXAMPLE IV

	<u>mg/ml</u>
30 Compound Ia	10
Methylcellulose	5.0
Tween 80	0.5
35 Benzyl alcohol	9.0
Benzalkonium chloride	1.0
Water to 1 ml.	

General Information

For Echinocandin B

45 *Emericella nidulans* (a later growth stage of *Aspergillus nidulans*, established as a producing organism for Echinocandin B) and maintained as MF 5100 in the Merck culture collection was employed in the fermentation. A culture was deposited with the American Type Culture Collection under the Budapest Treaty and assigned ATCC No. 20956.

50 Echinocandin B, prepared as described in Example C was compared for percent purity against previously prepared Echinocandin B having the following MS and NMR spectra and identified by comparison with published structure for Echinocandin B.

Mass Spectral Data: FAB mass spectra were recorded on a VG 20-253 mass spectrometer. A matrix of dithiothreitol-dithiocrythritol-Lil was used and M+Li was observed at m/z 1066 corresponding to the molecular weight 1059.

55 ¹H NMR Spectrum: The spectrum, Fig. 2, was recorded in CD₃OD on a Varian XL-300 NMR Spectrometer at 300 MHz. Chemical shifts are shown in ppm relative to tetramethylsilane (TMS) at zero ppm using the solvent peak at 3.30 ppm as internal reference.

¹³C NMR Data: The spectrum was recorded in CD₃OD at 24°C on a Varian XL-400 NMR spectrometer at 100 MHz. Chemical shifts are given in ppm relative to tetramethylsilane at zero ppm using the solvent peak at 49.0 ppm as internal reference.

¹³C Chemical Shifts (CD₃OD, 100 MHz): 11.28, 14.43, 19.66, 20.04, 23.62, 26.54, 27.05, 28.15, 28.20, 30.27(x2), 30.42, 30.46, 30.78, 32.65, 35.08, 36.82, 38.60, 39.06, 51.67, 52.94, 56.32, 56.84, 57.19, 58.61, 62.46, 68.33, 69.59, 69.70, 70.87, 71.29, 74.60, 75.52, 75.77, 76.94, 116.21(x2), 129.06(x2), 129.60(x2), 130.94, 130.96, 133.09, 158.46, 169.97, 172.48, 172.51, 172.79, 173.53, 174.35, 176.19.

10 Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. The use of a cyclohexapeptide compound represented by the formula


15

20

25

30

35




wherein

R' is H or OH;

R'' is H or OH;

R''' is H, OH, -O-alkyl(C₁-C₆), -O-benzyl, or

40



45

wherein R¹ is H, C₁-C₁₂ alkyl, C₃-C₇ cycloalkyl, hydroxyethyl, C₁₋₆ alkoxyethyl, furylmethyl, tetrahydrofurylmethyl, phenyl, phenylalkyl wherein the phenyl may be substituted with halo, hydroxy, C₁₋₆ alkyl and C₁₋₆ alkoxy;

50

R² is hydrogen or C₁₋₆ alkyl; or R¹ and R² together form -(CH₂)₂Q(CH₂)₂- wherein Q is -CH₂-, -NH- or -N-(C₁₋₆)-alkyl;

and m is from 0 to 4;

R''' is H or OH;


R'''' is H, CH₃ or CH₂CONH₂;

55


R'''' is H or CH₃; and

R is


(A)




from 6 to 24 carbon atoms
(B)




from 6 to 24 carbon atoms
(C) when R', R'', R''' and R'''' are OH, R'''' is CH₃ then
(i)



(ii)



or
(iii)



wherein A is divalent oxygen, sulfur, sulfinyl or sulfonyl;

A' is divalent oxygen, sulfur, sulfinyl, sulfonyl or -NH-;

X is hydrogen, chloro, bromo, iodo, nitro, C₁-C₃ alkyl, hydroxy, C₁-C₃ alkoxy, mercapto, C₁-C₃ alkylthio, carbamyl or C₁-C₃ alkylcarbamyl;

X' is chloro, bromo or iodo,


R₁ is hydrogen, C₁-C₁₈alkyl or C₂-C₁₈alkenyl;

W is C₁-C₁₀alkylene or C₂-C₁₀alkenylene;


m, n and p are 0 or 1, but if m is 0, n must be 0;

provided that the sum of the carbon atoms in the R₁ and W groups must be greater than 4 but cannot exceed 21; that when X is mercapto, A and A' cannot be sulfinyl or sulfonyl; and that when A and A' are sulfinyl or sulfonyl, they must be in equal oxidation states;


(D) when R' is OH, and R'', R''' and R'''' are H, then




wherein Y is a divalent aminoacyl radical of the formula
a)



wherein Z is C₁-C₁₀ alkylene or C₅-C₈ cycloalkylene
b)




wherein R₃ is hydroxymethyl, hydroxyethyl, mercaptomethyl, mercaptoethyl, methylthioethyl, 2-thienyl, 3-indolomethyl, phenyl, benzyl, halophenyl, halobenzyl, C₁-C₃ alkylphenyl, C₁-C₃ alkylbenzyl, C₁-C₃ alkylthiophenyl, C₁-C₃ alkylthiobenzyl, carbamylphenyl, carbamylbenzyl, C₁-C₃ alkylcarbamylphenyl or C₁-C₃ alkylcarbamylbenzyl;
(c)



wherein X² is hydrogen or halo; for the preparation of a medicament, useful for the treatment of or for the prevention of Pneumocystis carinii infections.

2. The use according to Claim 1 wherein the cyclohexapeptide compound is administered at a dose of from 0.5 to 20.0 mg/kg of mammalian body weight.
3. The use according to Claim 1 wherein the cyclohexapeptide compound is administered to a patient with an impaired immune system infected with Pneumocystis carinii.
4. The use according to Claim 1 wherein the cyclohexapeptide compound is administered to a patient infected with Pneumocystis carinii and exhibits AIDS.
5. A composition useful for the treatment of Pneumocystis carinii infections in mammals and adapted for administration by inhalation, comprising a pharmaceutically acceptable carrier and a cyclohexapeptide compound having the formula




wherein

R' is H or OH;

R'' is H or OH;

R''' is H, OH, -O-alkyl(C₁-C₆), -O-benzyl, or



wherein R¹ is H, C₁-C₁₂ alkyl, C₃-C₇ cycloalkyl, hydroxyethyl, C₁₋₆ alkoxyethyl, furylmethyl, tetrahydrofurylmethyl, phenyl, phenylalkyl wherein the phenyl may be substituted with halo, hydroxy, C₁₋₆ alkyl and C₁₋₆ alkoxy; R² is hydrogen or C₁₋₆ alkyl; or R¹ and R² together form -(CH₂)₂Q(CH₂)₂- wherein Q is -CH₂-, -NH- or -N-(C₁₋₆)-alkyl; and m is from 0 to 4;


R''' is H or OH;

R'''' is H, CH₃ or CH₂CONH₂; and

R'''' is H or CH₃; and


R is

(A)



from 6 to 24 carbon atoms


(B)




from 6 to 24 carbon atoms

(C) when R', R'', R''' and R'''' are OH, R'''' is CH₃ then

(i)




(ii)



or

(iii)



wherein A is divalent oxygen, sulfur, sulfinyl or sulfonyl;

A¹ is divalent oxygen, sulfur, sulfinyl, sulfonyl or -NH-;

X is hydrogen, chloro, bromo, iodo, nitro, C₁-C₃ alkyl, hydroxy, C₁-C₃ alkoxy, mercapto, C₁-C₃ alkylthio, carbamyl or C₁-C₃ alkylcarbamyl;

X¹ is chloro, bromo or iodo,


R¹ is hydrogen, C₁-C₁₈ alkyl or C₂-C₁₈ alkenyl;

W is C₁-C₁₀alkylene or C₂-C₁₀ alkenylene;

m, n and p are 0 or 1, but if m is 0, n must be 0;


provided that the sum of the carbon atoms in the R₁ and W groups must be greater than 4 but cannot exceed 21; that when X is mercapto A and A¹ cannot be sulfinyl or sulfonyl; and that when A and A¹ are sulfinyl or sulfonyl, they must be in equal oxidation states;

(D) when R' is OH, and R'', R''' and R'''' are H, then




wherein Y is a divalent aminoacyl radical of the formula

a)




wherein Z is C₁-C₁₀ alkylene or C₅-C₆ cycloalkylene

b)




wherein R₃ is hydroxyethyl, hydroxyethyl, mercaptomethyl, mercaptoethyl, methylthioethyl, 2-thienyl, 3-indolomethyl, phenyl, benzyl, halophenyl, halobenzyl, C₁-C₃ alkylphenyl, C₁-C₃ alkylbenzyl, C₁-C₃ alkylthiophenyl, C₁-C₃ alkylthiobenzyl, carbamylphenyl, carbamylbenzyl, C₁-C₃ alkylcarbamylphenyl or C₁-C₃ alkylcarbamylbenzyl;

(c)



wherein X² is hydrogen or halo.


6. The use according to Claim 1 wherein the compound is that having the formula



- 20
7. The use according to Claim 1 wherein the compound is 1-[(4R,5R)-4,5-dihydroxy-N²-(10,12-dimethyl-1-oxotetradecyl)-L-ornithine-5-(threo-3-hydroxy-1-glutamine)echinocandin B.
- 25
8. The use, according to Claim 1 wherein the compound is 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithine]echinocandin B.
9. The use according to Claim 1 wherein the compound is echinocandin B.
10. The use according to Claim 1 wherein the compound is tetrahydroechinocandin B.
- 30
11. The use according to Claim 1 wherein the compound is O-methyltetrahydroechinocandin B.
12. The use according to Claim 1 wherein the compound is O-benzyltetrahydroechinocandin B.

Claims for the following Contracting State : ES

- 35
1. A process for the preparation of a composition useful for the treatment of *Pneumocystis carinii* infections in mammals and adapted for administration by inhalation, which comprises admixing a pharmaceutically acceptable carrier with a cyclohexapeptide compound having the formula
- 40
- 45
- 50
- 55




wherein

R' is H or OH;

R'' is H or OH;

R''' is H, OH, -O-alkyl(C₁-C₆), -O-benzyl, or



wherein R' is H, C₁-C₁₂ alkyl, C₃-C₇ cycloalkyl, hydroxyethyl, C₁₋₆ alkoxyethyl, furylmethyl, tetrahydrofurylmethyl, phenyl, phenylalkyl wherein the phenyl may be substituted with halo, hydroxy, C₁₋₆ alkyl and C₁₋₆ alkoxy; R² is hydrogen or C₁₋₆ alkyl; or R¹ and R² together form -(CH₂)₂Q(CH₂)₂- wherein Q is -CH₂-, -NH- or -N-(C₁₋₆)-alkyl; and m is from 0 to 4;


R''' is H or OH;

R'''' is H, CH₃ or CH₂CONH₂; and

R''''' is H or CH₃; and


R is

(A)



from 6 to 24 carbon atoms


(B)





from 6 to 24 carbon atoms

(C) when R', R'', R''' and R'''' are OH R''''' is CH₃ then

(i)



(ii)

or
(iii)

wherein A is divalent oxygen, sulfur, sulfinyl or sulfonyl;

A' is divalent oxygen, sulfur, sulfinyl, sulfonyl or -NH-;

X is hydrogen, chloro, bromo, iodo, nitro, C₁-C₃ alkyl, hydroxy, C₁-C₃ alkoxy, mercapto, C₁-C₃ alkylthio, carbamyl or C₁-C₃ alkylcarbamyl;

X' is chloro, bromo or iodo,

R₁ is hydrogen, C₁-C₁₈ alkyl or C₂-C₁₈ alkenyl;W is C₁-C₁₀ alkylene or C₂-C₁₀ alkenylene;


m, n and p are 0 or 1, but if m is 0, n must be 0;

provided that the sum of the carbon atoms in the R₁ and W groups must be greater than 4

but cannot exceed 21; that when X is mercapto A and A' cannot be sulfinyl or sulfonyl; and that


when A and A' are sulfinyl or sulfonyl, they must be in equal oxidation states;

(D) when R' is OH, and R'', R''' and R'''' are H, then




wherein Y is a divalent aminoacyl radical of the formula


a)

wherein Z is C₁-C₁₀ alkylene or C₅-C₆ cycloalkylene

b)


wherein R₃ is hydroxyethyl, hydroxyethyl, mercaptomethyl, mercaptoethyl, methylthioethyl, 2-thienyl, 3-indolomethyl, phenyl, benzyl, halophenyl, halobenzyl, C₁-C₃ alkylphenyl, C₁-C₃ alkylbenzyl, C₁-C₃ alkylthiophenyl, C₁-C₃ alkylthiobenzyl, carbamylphenyl, carbamylbenzyl, C₁-C₃ alkylcarba-

mylphenyl or C₁-C₃ alkylcarbonylbenzyl;
(c)



wherein X² is hydrogen or halo.

2. A process according to Claim 1 wherein the compound is that having the formula




3. A process according to Claim 1 wherein the compound is 1-[(4R,5R)-4,5-dihydroxy-N²-(10,12-dimethyl-1-oxotetradecyl)-L-ornithine-5-(threo-3-hydroxy-1-glutamine)echinocandin B.
4. A process according to Claim 1 wherein the compound is 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithine]echinocandin B.
5. A process according to Claim 1 wherein the compound is echinocandin B.
6. A process according to Claim 1 wherein the compound is tetrahydroechinocandin B.
7. A process according to Claim 1 wherein the compound is O-methyltetrahydroechinocandin B.
8. A process according to Claim 1 wherein the compound is O-benzyltetrahydroechinocandin B.

Patentansprüche


Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Verwendung einer Cyclohexapeptidverbindung, dargestellt durch die Formel


(i)



(ii)



oder
(iii)



in denen A bivalenten Sauerstoff, Schwefel, Sulfinyl oder Sulfonyl;

A' bivalenten Sauerstoff, Schwefel, Sulfinyl, Sulfonyl oder -NH-;

X Wasserstoff, Chlor, Brom, Jod, Nitro, C₁-C₃ Alkyl, Hydroxy, C₁-C₃ Alkoxy, Mercapto, C₁-C₃ Alkylthio, Carbamoyl oder C₁-C₃ Alkylcarbamoyl;

X' Chlor, Brom oder Jod;


R' Wasserstoff, C₁-C₁₈ Alkyl oder C₂-C₁₈ Alkenyl;

und W C₁-C₁₀ Alkylen oder C₂-C₁₀ Alkenylen darstellen und

m, n und p 0 oder 1 sind, wobei jedoch, wenn m 0 ist, n auch 0 sein muß;


vorausgesetzt daß die Summe der Kohlenstoffatome in den Resten R₁ und W größer als 4 ist und 21 nicht überschreitet, daß A und A' nicht Sulfinyl oder Sulfonyl sein können, wenn X Mercapto bedeutet, und daß, wenn A und A' Sulfinyl oder Sulfonyl bedeuten, sie auf der gleichen Oxidationsstufe sein müssen;

(D) falls R' OH ist und R'', R''' und R'''' H sind, dann




wobei Y ein bivalentes Aminoacylradikal der Formeln

a)




worin Z C₁-C₁₀ Alkylen oder C₅-C₈ Cycloalkylen ist,

b)




worin R_3 Hydroxymethyl, Hydroxyethyl, Mercaptomethyl, Mercaptoethyl, Methylthioethyl, 2-Thienyl, 3-Indolmethyl, Phenyl, Benzyl, Halogenphenyl, Halogenbenzyl, C_1 - C_3 Alkylphenyl, C_1 - C_3 Alkylbenzyl, C_1 - C_3 Alkylthiophenyl, C_1 - C_3 Alkylthiobenzyl, Carbamoylphenyl, Carbamoylbenzyl, C_1 - C_3 Alkylcarbamoylphenyl oder C_1 - C_3 Alkylcarbamoylbenzyl ist; und
c)



worin X^2 Wasserstoff oder Halogen ist; darstellt,
für die Herstellung eines Arzneimittels, welches für die Behandlung oder für die Vorbeugung von Infektionen durch *Pneumocystis carinii* verwendbar ist.


2. Verwendung gemäß Anspruch 1, wobei die Cyclohexapeptidverbindung in einer Dosis von 0,5 bis 20,0 mg/kg Säugetier verabreicht wird.
3. Verwendung gemäß Anspruch 1, wobei die Cyclohexapeptidverbindung einem mit *Pneumocystis carinii* infizierten Patienten mit geschwächtem Immunsystem verabreicht wird.
4. Verwendung gemäß Anspruch 1, wobei die Cyclohexapeptidverbindung einem mit *Pneumocystis carinii* infizierten, an AIDS leidenden Patienten verabreicht wird.
5. Eine Zusammensetzung, die für die Behandlung von Infektionen durch *Pneumocystis carinii* bei Säugtieren verwendbar und für die Verabreichung durch Inhalation angepaßt ist, umfassend einen pharmazeutisch annehmbaren Träger und eine Cyclohexapeptidverbindung mit der Formel



(I)

in der

R' H oder OH;
 R'' H oder OH;
 R''' H, OH, -O-Alkyl(C_1 - C_6), -O-Benzyl oder



wobei R^1 H, C_1 - C_{12} Alkyl, C_3 - C_7 Cycloalkyl, Hydroxyethyl, C_{1-6} Alkoxyalkyl, Furylmethyl, Tetrahydrofurylmethyl, Phenyl, Phenylalkyl, worin der Phenylrest durch Halogen, Hydroxy, Cl-6 Alkyl oder C_{1-6} Alkoxy substituiert sein kann,

R^2 H oder C_{1-6} Alkyl oder


R^1 und R^2 zusammen den Rest $-(CH_2)_2Q(CH_2)_2-$, in dem Q $-CH_2-$, $-NH-$ oder $-N-(C_{1-6})$ Alkyl ist, darstellen und m eine ganze Zahl von 0 bis 4 ist;

R''' H oder OH;

R'''' H, CH_3 oder CH_2CONH_2 ;


R''''' H oder CH_3 bedeuten und in der R eine der folgenden Bedeutungen aufweist:

(A)



mit 6 bis 24 Kohlenstoffatomen,


(B)




mit 6 bis 24 Kohlenstoffatomen,

(C) falls R' , R'' , R''' und R'''' OH sind und R''''' CH_3 ist, einen der Reste

(i)




(ii)



oder

(iii)



in denen A bivalenten Sauerstoff, Schwefel, Sulfinyl oder Sulfonyl;

A' bivalenten Sauerstoff, Schwefel, Sulfinyl, Sulfonyl oder -NH-;

X Wasserstoff, Chlor, Brom, Jod, Nitro, C₁-C₃ Alkyl, Hydroxy, C₁-C₃ Alkoxy, Mercapto, C₁-C₃ Alkylthio, Carbamoyl oder C₁-C₃ Alkylcarbamoyl;

X' Chlor, Brom oder Jod;

R' Wasserstoff, C₁-C₁₈ Alkyl oder C₂-C₁₈ Alkenyl; und W C₁-C₁₀ Alkylen oder C₂-C₁₀ Alkenylen darstellen und

m, n und p 0 oder 1 sind, wobei jedoch, wenn m 0 ist, n auch 0 sein muß;

vorausgesetzt daß die Summe der Kohlenstoffatome in den Resten R' und W größer als 4 ist und 21 nicht überschreitet, daß A und A' nicht Sulfinyl oder Sulfonyl sein können, wenn X Mercapto bedeutet, und daß, wenn A und A' Sulfinyl oder Sulfonyl bedeuten, sie auf der gleichen Oxidationsstufe sein müssen;

(D) falls R' OH ist und R'', R''' und R'''' H sind, dann



wobei Y ein bivalentes Aminoacylradikal der Formeln

a)



35 worin Z C₁-C₁₀ Alkylen oder C₅-C₈ Cycloalkylen ist,

b)




45 worin R₃ Hydroxymethyl, Hydroxyethyl, Mercaptomethyl, Mercaptoethyl, Methylthioethyl, 2-Thienyl, 3-Indolmethyl, Phenyl, Benzyl, Halogenphenyl, Halogenbenzyl, C₁-C₃ Alkylphenyl, C₁-C₃ Alkylbenzyl, C₁-C₃ Alkylthiophenyl, C₁-C₃ Alkylthiobenzyl, Carbamoylphenyl, Carbamoylbenzyl, C₁-C₃ Alkylcarbamoylphenyl oder C₁-C₃ Alkylcarbamoylbenzyl ist; und

c)



55 worin X² Wasserstoff oder Halogen ist; darstellt.

6. Verwendung gemäß Anspruch 1, wobei es sich um eine Verbindung mit der Formel




handelt.

7. Verwendung gemäß Anspruch 1, wobei die Verbindung 1-[(4R,5R)-4,5-Dihydroxy-N²-(10,12-dimethyl-1-oxotetradecyl)-L-ornithin]-5-(threo-3-hydroxy-1-glutamin)echinocandin B ist.
8. Verwendung gemäß Anspruch 1, wobei die Verbindung 1-[(4R,5R)-4,5-Dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithin]-echinocandin B ist.
9. Verwendung gemäß Anspruch 1, wobei die Verbindung Echinocandin B ist.
10. Verwendung gemäß Anspruch 1, wobei die Verbindung Tetrahydroechinocandin B ist.
11. Verwendung gemäß Anspruch 1, wobei die Verbindung O-Methyltetrahydroechinocandin B ist.
12. Verwendung gemäß Anspruch 1, wobei die Verbindung O-Benzyltetrahydroechinocandin B ist.

Patentansprüche für folgende Vertragsstaat : ES

1. Verfahren für die Herstellung einer Zusammensetzung, die für die Behandlung von Infektionen durch *Pneumocystis carinii* verwendbar und für die Verabreichung mittels Inhalation angepaßt ist, umfassend die Mischung eines pharmazeutisch annehmbaren Trägers mit einer Cyclohexapeptidverbindung der Formel




in der

R' H oder OH;

R'' H oder OH;

R''' H, OH, -O-Alkyl(C_1 - C_6), -O-Benzyl oder



wobei R^1 H, C_1 - C_{12} Alkyl, C_3 - C_7 Cycloalkyl, Hydroxyethyl, C_{1-6} Alkoxyalkyl, Furylmethyl, Tetrahydrofurylmethyl, Phenyl, Phenylalkyl, worin der Phenylrest durch Halogen, Hydroxy, Cl-6 Alkyl oder C_{1-6} Alkoxy substituiert sein kann,

R^2 H oder C_{1-6} Alkyl oder

R^1 und R^2 zusammen den Rest $-(\text{CH}_2)_2\text{Q}(\text{CH}_2)_2-$, in dem Q $-\text{CH}_2-$, $-\text{NH}-$ oder $-\text{N}-(\text{C}_{1-6})\text{Alkyl}$ ist, darstellen und m 0 bis 4 ist;


R''' H oder OH;

R'''' H, CH_3 oder CH_2CONH_2 ;

R''''' H oder CH_3 bedeuten und in der


R eine der folgenden Bedeutungen aufweist:

(A)




mit 6 bis 24 Kohlenstoffatomen,

(B)




mit 6 bis 24 Kohlenstoffatomen,

(C) falls R', R'', R''' und R'''' OH sind und R'''' CH₃ ist, einen der Reste
(i)




(ii)



oder

(iii)



in denen A bivalenten Sauerstoff, Schwefel, Sulfinyl oder Sulfonyl;

A' bivalenten Sauerstoff, Schwefel, Sulfinyl, Sulfonyl oder -NH-;

X Wasserstoff, Chlor, Brom, Jod, Nitro, C₁-C₃ Alkyl, Hydroxy, C₁-C₃ Alkoxy, Mercapto, C₁-C₃ Alkylthio, Carbamoyl oder C₁-C₃ Alkylcarbamoyl;

X' Chlor, Brom oder Jod;


R₁ Wasserstoff, C₁-C₁₈ Alkyl oder C₂-C₁₈ Alkenyl;

und W C₁-C₁₀ Alkylen oder C₂-C₁₈ Alkenylen darstellen und

m, n und p 0 oder 1 sind, wobei jedoch, wenn m 0 ist, n auch 0 sein muß;


vorausgesetzt daß die Summe der Kohlenstoffatome in den Resten R₁ und W größer als 4 ist und 21 nicht überschreitet, daß A und A' nicht Sulfinyl oder Sulfonyl sein können, wenn X Mercapto bedeutet, und daß, wenn A und A' Sulfinyl oder Sulfonyl bedeuten, sie auf der gleichen Oxidationsstufe sein müssen;

(D) falls R' OH ist und R'', R''' und R'''' H sind, dann




wobei Y ein bivalentes Aminoacylradikal der Formeln

a)




worin Z C₁-C₁₀ Alkylen oder C₅-C₈ Cycloalkylen ist,

b)




worin R₃ Hydroxymethyl, Hydroxyethyl, Mercaptomethyl, Mercaptoethyl, Methylthioethyl, 2-Thienyl, 3-Indolylmethyl, Phenyl, Benzyl, Halogenphenyl, Halogenbenzyl, C₁-C₃ Alkylphenyl, C₁-C₃ Alkylbenzyl, C₁-C₃ Alkylthiophenyl, C₁-C₃ Alkylthiobenzyl, Carbamoylphenyl, Carbamoylbenzyl, C₁-C₃ Alkylcarbamoylphenyl oder C₁-C₃ Alkylcarbamoylbenzyl ist; und
c)



worin X² Wasserstoff oder Halogen ist; darstellt.

2. Verfahren gemäß Anspruch 1, wobei es sich um eine Verbindung der Formel




handelt.

3. Verfahren gemäß Anspruch 1, wobei die Verbindung 1-[(4R,5R)-4,5-Dihydroxy-N²-(10,12-dimethyl-1-oxotetradecyl)-L-ornithin]-5-(threo-3-hydroxy-1-glutamin)-echinocandin B ist.
4. Verfahren gemäß Anspruch 1, wobei die Verbindung 1-[(4R,5R)-4,5-Dihydroxy-N²-(4-(octyloxy)benzoyl)-L-ornithin]-echinocandin B ist.
5. Verfahren gemäß Anspruch 1, wobei die Verbindung Echinocandin B ist.
6. Verfahren gemäß Anspruch 1, wobei die Verbindung Tetrahydroechinocandin B ist.
7. Verfahren gemäß Anspruch 1, wobei die Verbindung O-Methyltetrahydroechinocandin B ist.
8. Verfahren gemäß Anspruch 1, wobei die Verbindung O-Benzyltetrahydroechinocandin B ist.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Utilisation d'un composé cyclohexapeptidique représenté par la formule




dans laquelle

R' est H ou OH ;

R'' est H ou OH ;

R''' est H, OH, -O-alkyle(C_1 - C_6), -O-benzyle ou



dans laquelle R^1 est H, un alkyle en C_1 - C_{12} , un cycloalkyle en C_3 - C_7 , un hydroxyéthyle, un (alcoxy en C_{1-6})éthyle, un furylméthyle, un tétrahydrofurylméthyle, un phényle, un phénylalkyle dont le phényle peut être substitué avec un halogéno, un hydroxy, un alkyle en C_{1-6} et un alcoxy en C_{1-6} ; R^2 est un hydrogène ou un alkyle en C_{1-6} ; ou R^1 et R^2 forment ensemble $-(CH_2)_2Q(CH_2)_2-$ où Q est $-CH_2-$, $-NH-$ ou $-N$ -alkyl(C_{1-6}) ; et m a une valeur de 0 à 4 ;


R''' est H ou OH ;

R'''' est H, CH_3 ou CH_2CONH_2 ; et

R'''' est H ou CH_3


R est

(A)



de 6 à 24 atomes de carbone,


(B)




de 6 à 24 atomes de carbone,

(C), lorsque R', R'', R''' et R'''' sont OH et R'''' est CH₃, il est

(i)




(ii)



ou

(iii)



où

A est un oxygène, un soufre, un sulfinyle ou un sulfonyle divalents ;

A' est un oxygène, un soufre, un sulfinyle ou un sulfonyle divalents ou -NH- ;

X est un hydrogène, un chloro, un bromo, un iodo, un nitro, un alkyle en C₁-C₃, un hydroxy, un alcoxy en C₁-C₃, un mercapto, un alkylthio en C₁-C₃, un carbamyle ou un (alkyl en C₁-C₃)carbamyle ;

X' est un chloro, un bromo ou un iodo ;

R' est un hydrogène, un alkyle en C₁-C₁₈ ou un alcényle en C₂-C₁₈

W est un alkylène en C₁-C₁₀ ou un alcénylène en C₂-C₁₀ ;

m, n et p sont 0 ou 1, mais si m est 0, n doit être 0 ;


sous réserve que la somme des atomes de carbone dans les groupes R' et W soit supérieure à 4 mais ne puisse pas dépasser 21 ; que, lorsque X est un mercapto, A et A' ne puissent pas être un sulfinyle ou un sulfonyle ; et que, lorsque A et A' sont un sulfinyle ou un sulfonyle, ils soient à des états d'oxydation égaux ;

(D), lorsque R' est OH, R'', R''' et R'''' sont H, il est



où Y est un radical aminoacycle divalent de formule

a)



où Z est un alkylène en C₁-C₁₀ ou un cycloalkylène en C₅-C₈

b)




où R_3 est un hydroxyméthyle, un hydroxyéthyle, un mercaptométhyle, un mercaptoéthyle, un méthylthioéthyle, un 2-thiényle, un 3-indolométhyle, un phényle, un benzyle, un halogénophényle, un halogénobenzyle, un (alkyl en C_1-C_3) phényle, un (alkyl en C_1-C_3) benzyle, un (alkyl en C_1-C_3) thio-phényle, un (alkyl en C_1-C_3) thiobenzyle, un carbamylphényle, un carbamylbenzyle, un (alkyl en C_1-C_3) carbamylphényle ou un (alkyl en C_1-C_3) carbamylbenzyle ;

c)



20 où X^2 est un hydrogène ou un halogéno ; pour la préparation d'un médicament utile pour le traitement ou pour la prévention des infections à Pneumocystis carinii.

2. Utilisation selon la revendication 1, dans laquelle le composé cyclohexapeptidique est administré à une dose de 0,5 à 20,0 mg/kg de poids corporel de mammifère.
3. Utilisation selon la revendication 1, dans laquelle le composé cyclohexapeptidique est administré à un patient dont le système immunitaire est altéré infecté par Pneumocystis carinii.
4. Utilisation selon la revendication 1, dans laquelle le composé cyclohexapeptidique est administré à un patient infecté par Pneumocystis carinii et atteint de SIDA.
5. Composition utile pour le traitement des infections à Pneumocystis carinii des mammifères et adaptée à l'administration par inhalation, comprenant un véhicule pharmaceutiquement acceptable et un composé cyclohexapeptidique de formule




55 dans laquelle

R' est H ou OH ;

R'' est H ou OH ;

R''' est H, OH, -O-alkyle(C_1-C_6), -O-benzyle ou



dans laquelle R¹ est H, un alkyle en C₁-C₁₂, un cycloalkyle en C₃-C₇, un hydroxyéthyle, un (alcoxy en C₁₋₆)éthyle, un furylméthyle, un tétrahydrofurylméthyle, un phényle, un phénylalkyle dont le phényle peut être substitué avec un halogéno, un hydroxy, un alkyle en C₁₋₆ et un alcoxy en C₁₋₆ ; R² est un hydrogène ou un alkyle en C₁₋₆ ; ou R¹ et R² forment ensemble -(CH₂)₂Q(CH₂)₂- où Q est -CH₂-, -NH- ou -N-alkyl(C₁₋₆)- ; et m a une valeur de 0 à 4 ;


R''' est H ou OH ;

R'''' est H, CH₃ ou CH₂CONH₂ ; et

R'''''' est H ou CH₃


R est

(A)



de 6 à 24 atomes de carbone,


(B)




de 6 à 24 atomes de carbone,

(C), lorsque R', R'', R''' et R'''' sont OH et R'''''' est CH₃, il est

(i)




(ii)



ou

(iii)



où

A est un oxygène, un soufre, un sulfinyle ou un sulfonyle divalents ;

A¹ est un oxygène, un soufre, un sulfinyle ou un sulfonyle divalents ou -NH- ;

X est un hydrogène, un chloro, un bromo, un iodo, un nitro, un alkyle en C₁-C₃, un hydroxy, un alcoxy en C₁-C₃, un mercapto, un alkylthio en C₁-C₃, un carbamyle ou un (alkyl en C₁-C₃)carbamyle ;

X¹ est un chloro, un bromo ou un iodo ;

5 R¹ est un hydrogène, un alkyle en C₁-C₁₈ ou un alcényle en C₂-C₁₈ ;


W est un alkylène en C₁-C₁₀ ou un alcénylène en C₂-C₁₀ ;

m, n et p sont 0 ou 1, mais si m est 0, n doit être 0 ;

sous réserve que la somme des atomes de carbone dans les groupes R¹ et W soit supérieure à 4 mais ne puisse pas dépasser 21 ; que, lorsque X est un mercapto, A et A¹ ne puissent pas être un sulfinyle ou un sulfonyle ;

10 et que, lorsque A et A¹ sont un sulfinyle ou un sulfonyle, ils soient à des états d'oxydation égaux ;

(D), lorsque R' est OH, R'', R''' et R'''' sont H, il est



où Y est un radical aminoacyle divalent de formule

a)



où Z est un alkylène en C₁-C₁₀ ou un cycloalkylène en C₅-C₆

b)



30 où R₃ est un hydroxyméthyle, un hydroxyéthyle, un mercaptométhyle, un mercaptoéthyle, un méthylthioéthyle, un 2-thiényle, un 3-indolométhyle, un phényle, un benzyle, un halogénophényle, un halogénobenzyle, un (alkyl en C₁-C₃) phényle, un (alkyl en C₁-C₃)benzyle, un (alkyl en C₁-C₃) thio-phényle, un (alkyl en C₁-C₃)thiobenzyle, un carbamylphényle, un carbamylbenzyle, un (alkyl en C₁-C₃) carbamylphényle ou un (alkyl en C₁-C₃)carbamylbenzyle ;

35 c)




où X² est un hydrogène ou un halogéno.

6. Utilisation selon la revendication 1, dans laquelle le composé répond à la formule

45

50


55



- 20 7. Utilisation selon la revendication 1, dans laquelle le composé est la 1-[(4R,5R)-4,5-dihydroxy-N²-(10,12-diméthyl-1-oxotétradécyl)-L-ornithine-5-(thréo-3-hydroxy-1-glutamine)] échinocandine B.
8. Utilisation selon la revendication 1, dans laquelle le composé est la 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithine]échinocandine B.
- 25 9. Utilisation selon la revendication 1, dans laquelle le composé est l'échinocandine B.
10. Utilisation selon la revendication 1, dans laquelle le composé est la tétrahydroéchinocandine B.
11. Utilisation selon la revendication 1, dans laquelle le composé est l'O-méthyltétrahydroéchinocandine B.
- 30 12. Utilisation selon la revendication 1, dans laquelle le composé est l'O-benzyltétrahydroéchinocandine B.

Revendications pour l'Etat contractant suivant : ES

- 35 1. Procédé pour la préparation d'une composition utile pour le traitement des infections à *Pneumocystis carinii* des mammifères et adaptée à l'administration par inhalation, qui comprend le mélange d'un véhicule pharmaceutiquement acceptable avec un composé cyclohexapeptidique de formule




dans laquelle

R' est H ou OH ;

R'' est H ou OH ;

R''' est H, OH, -O-alkyle(C₁-C₆), -O-benzyle ou



dans laquelle R¹ est H, un alkyle en C₁-C₁₂, un cycloalkyle en C₃-C₇, un hydroxyéthyle, un (alcoxy en C₁₋₆)éthyle, un furylméthyle, un tétrahydrofurylméthyle, un phényle, un phénylalkyle dont le phényle peut être substitué avec un halogéno, un hydroxy, un alkyle en C₁₋₆ et un alcoxy en C₁₋₆ ; R² est un hydrogène ou un alkyle en C₁₋₆ ; ou R¹ et R² forment ensemble -(CH₂)₂Q(CH₂)₂- où Q est -CH₂-, -NH- ou -N-alkyl(C₁₋₆)- ; et m a une valeur de 0 à 4 ;


R''' est H ou OH ;

R'''' est H, CH₃ ou CH₂ CONH₂ ; et

R'''' est H ou CH₃ ; et


R est

(A)



de 6 à 24 atomes de carbone,


(B)




de 6 à 24 atomes de carbone,

(C), lorsque R', R'', R''' et R'''' sont OH et R'''' est CH₃, il est

(i)




(ii)



ou

(iii)



où

A est un oxygène, un soufre, un sulfinyle ou un sulfonyle divalents ;

A' est un oxygène, un soufre, un sulfinyle ou un sulfonyle divalents ou -NH- ;

X est un hydrogène, un chloro, un bromo, un iodo, un nitro, un alkyle en C₁-C₃, un hydroxy, un alcoxy en C₁-C₃, un mercapto, un alkylthio en C₁-C₃, un carbamyle ou un (alkyl en C₁-C₃)carbamyle ;

X' est un chloro, un bromo ou un iodo ;


R' est un hydrogène, un alkyle en C₁-C₁₈ ou un alcényle en C₂-C₁₈ ;

W est un alkylène en C₁-C₁₀ ou un alcénylène en C₂-C₁₀ ;

m, n et p sont 0 ou 1, mais si m est 0, n doit être 0 ;


sous réserve que la somme des atomes de carbone dans les groupes R' et W soit supérieure à 4 mais ne puisse pas dépasser 21 ; que, lorsque X est un mercapto, A et A' ne puissent pas être un sulfinyle ou un sulfonyle ; et que, lorsque A et A' sont un sulfinyle ou un sulfonyle, ils soient à des états d'oxydation égaux ;

(D), lorsque R' est OH, R'', R''' et R'''' sont H, il est




où Y est un radical aminoacyle divalent de formule

a)




où Z est un alkylène en C₁-C₁₀ ou un cycloalkylène en C₅-C₆

b)




où R₃ est un hydroxyméthyle, un hydroxyéthyle, un mercaptométhyle, un mercaptoéthyle, un méthylthioéthyle, un 2-thiényle, un 3-indolométhyle, un phényle, un benzyle, un halogénophényle, un halogénobenzyle, un (alkyl en C₁-C₃) phényle, un (alkyl en C₁-C₃) benzyle, un (alkyl en C₁-C₃) thio-phényle, un (alkyl en C₁-C₃)thiobenzyle, un carbamylphényle, un carbamylbenzyle, un (alkyl en C₁-C₃) carbamylphényle ou un (alkyl en C₁-C₃)carbamylbenzyle ;

c)



où X² est un hydrogène ou un halogéno.

2. Procédé selon la revendication 1, dans lequel le composé a pour formule



3. Procédé selon la revendication 1, dans lequel le composé est la 1-[(4R,5R)-4,5-dihydroxy-N²-(10,12-diméthyl-1-oxo-tétradécyl)-L-ornithine-5-(thréo-3-hydroxy-1-glutamine)]échinocandine B.
4. Procédé selon la revendication 1, dans lequel le composé est la 1-[(4R,5R)-4,5-dihydroxy-N²-[4-(octyloxy)benzoyl]-L-ornithine]échinocandine B.
5. Procédé selon la revendication 1, dans lequel le composé est l'échinocandine B.
6. Procédé selon la revendication 1, dans lequel le composé est la tétrahydroéchinocandine B.
7. Procédé selon la revendication 1, dans lequel le composé est l'O-méthyltétrahydroéchinocandine B.
8. Procédé selon la revendication 1, dans lequel le composé est l'O-benzyltétrahydroéchinocandine B.

